

08/25/00  
JCS03 U.S. PTO

# BERESKIN & PARR

BOX 401, 40 KING STREET WEST, TORONTO, CANADA M5H 3Y2  
PHONE (416) 364-7311 • FAX (416) 361-1398 • WWW.BERESKINPARR.COM

PTO  
09/645593  
08/25/00

Please type a plus sign (+) inside this box → ☒

PTO/SB/05 (4/98)  
Approved for use through 09/30/2000. OMB 0651-0032  
Patent and Trademark Office: U.S. DEPARTMENT OF COMMERCE  
Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

<b>UTILITY PATENT APPLICATION TRANSMITTAL</b> <small>(Only for new nonprovisional applications under 37 C.F.R. § 1.53(b))</small>	Attorney Docket No.	9369-151/MG
	First Inventor or Application Identifier	Sarita Chaudhary
	Title	Flax Seed Specific Promoters
	Express Mail Label No.	

<b>APPLICATION ELEMENTS</b> <small>See MPEP chapter 600 concerning utility patent application contents.</small>	<b>ADDRESS TO:</b> Assistant Commissioner for Patents Box Patent Application Washington, DC 20231
--	---

<p>1. <input type="checkbox"/> * Fee Transmittal Form (e.g., PTO/SB/17) (Submit an original and a duplicate for fee processing)</p> <p>2. <input checked="" type="checkbox"/> Specification [Total Pages 52] (preferred arrangement set forth below)</p> <ul style="list-style-type: none"><li>- Descriptive title of the invention</li><li>- Cross References to Related Applications</li><li>- Statement Regarding Fed sponsored R &amp; D</li><li>- Reference to Micro fiche Appendix</li><li>- Background of the invention</li><li>- Brief Summary of the invention</li><li>- Brief Description of the Drawings (if filed)</li><li>- Detailed Description</li><li>- Claim(s)</li><li>- Abstract of the Disclosure</li></ul> <p>3. <input checked="" type="checkbox"/> Drawing(s) (35 U.S.C. 113) [Total Sheets 24]</p> <p>4. Oath or Declaration [Total Pages ]</p> <p>a. <input type="checkbox"/> Newly executed (original or copy)</p> <p>b. <input type="checkbox"/> Copy from a prior application (37 C.F.R. § 1.63(d)) (for continuation/divisional with Box 16 completed)</p> <p>i. <input type="checkbox"/> <b>DELETION OF INVENTOR(S)</b> Signed statement attached deleting inventor(s) named in the prior application, see 37 C.F.R. §§ 1.63(d)(2) and 1.33(b).</p>	<p>5. <input type="checkbox"/> Microfiche Computer Program (Appendix)</p> <p>6. Nucleotide and/or Amino Acid Sequence Submission (if applicable, all necessary)</p> <p>a. <input checked="" type="checkbox"/> Computer Readable Copy</p> <p>b. <input checked="" type="checkbox"/> Paper Copy (identical to computer copy)</p> <p>c. <input type="checkbox"/> Statement verifying identity of above copies</p>
---	--

**ACCOMPANYING APPLICATION PARTS**

7. ☐ Assignment Papers (cover sheet & document(s))

8. ☐ 37 C.F.R. § 3.73(b) Statement ☐ Power of Attorney  
(when there is an assignee)

9. ☐ English Translation Document (if applicable)

10. ☐ Information Disclosure Statement (IDS)/PTO-1449 ☐ Copies of IDS Citations

11. ☐ Preliminary Amendment

12. ☒ Return Receipt Postcard (MPEP 503)  
(Should be specifically itemized)

13. ☐ \* Small Entity Statement(s) ☐ Statement filed in prior application, Status still proper and desired  
(PTO/SB/09-12)

14. ☐ Certified Copy of Priority Document(s)  
(if foreign priority is claimed)

15. ☐ Other: \_\_\_\_\_

\*NOTE FOR ITEMS 1 & 13: IN ORDER TO BE ENTITLED TO PAY SMALL ENTITY FEES, A SMALL ENTITY STATEMENT IS REQUIRED (37 C.F.R. § 1.27), EXCEPT IF ONE FILED IN A PRIOR APPLICATION IS RELIED UPON (37 C.F.R. § 1.28).

16. If a CONTINUING APPLICATION, check appropriate box, and supply the requisite information below and in a preliminary amendment:

☐ Continuation ☐ Divisional ☐ Continuation-in-part (CIP) of prior application No. \_\_\_\_\_

Prior application information: Examiner \_\_\_\_\_ Group / Art Unit: \_\_\_\_\_

For CONTINUATION or DIVISIONAL APPS only: The entire disclosure of the prior application, from which an oath or declaration is supplied under Box 4b, is considered a part of the disclosure of the accompanying continuation or divisional application and is hereby incorporated by reference. The incorporation can only be relied upon when a portion has been inadvertently omitted from the submitted application parts.

**17. CORRESPONDENCE ADDRESS**

☒ Customer Number or Bar Code Label **1059** or ☐ Correspondence address below  
(Insert Customer No. or Attach bar code label here)

Name	Bereskin & Parr				
Address	Box 401 40 King Street West				
City	Toronto	State	Ontario	Zip Code	M5H 3Y2
Country	Canada	Telephone	(416) 364-7311	Fax	(416) 361-1398

Name (Print/Type)	Micheline Gravelle	Registration No. (Attorney/Agent)	40,261
Signature		Date	August 24, 2000

Burden Hour Statement: This form is estimated to take 0.2 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Office, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Box Patent Application, Washington, DC 20231.

B&P File No. 9369-151/MG

**BERESKIN & PARR**

**UNITED STATES**

005220-2654360

**Title:** Flax Seed Specific Promoters  
**Inventors:** Sarita Chaudhary, Gijs van Rooijen, Maurice Moloney  
and Surinder Singh

**Title:** Flax Seed Specific Promoters

This application claims benefit from United States provisional application no. 60/151,044, filed August 27, 1999 and United States  
5 provisional application no. 60/161,722 filed on October 27, 1999, both of which are incorporated herein by reference.

**FIELD OF THE INVENTION**

The present invention relates to plant genetic engineering methods useful for the alteration of the constituents of plant seeds. More  
10 specifically, the invention relates to promoters that have been obtained from flax and are capable of directing expression of non-native genes in flax seeds as well as the seeds of other plants.

**BACKGROUND OF THE INVENTION**

Flax or linseed (*Linum usitatissimum*) is a commercially  
15 important oilseed crop. Flax oil and meal are valuable raw materials derived from flax seed. A further economically significant raw material, flax fiber, is obtainable from the stem of the plant. The flax oil fraction is used for non-edible purposes, for example in the manufacture of varnish and paint, and has more recently become suited for use in the  
20 manufacture of a range of edible products, such as margarines and salad oils and dressings, by virtue of newly bred so called Linola cultivars (Green (1986) Can. J. Plant Sci, 66: 499-503). Flax meal is used primarily as a constituent of ruminant feeds while flax fibers are used in the manufacture of linen fabrics. Given its economic importance as a source  
25 for raw materials, it is desirable to further improve and diversify the available flax cultivar portfolio both with respect to agronomic performance, for example seed yield, resistance to pathogens and low climatic temperatures, and with respect to yield and quality of the raw materials to suit downstream applications. Although it is possible to  
30 obtain improved flax cultivars through conventional plant breeding, as evidenced by the development of the Linola cultivars, developing an elite agronomic plant line requires large investments in plant breeding due to

09645593 "082500  
005220" 2 6554960

the long timelines involved. Plant genetic engineering technology allows the isolation of genes directly from unrelated species and the transfer of these genes into elite agronomic backgrounds, thereby significantly reducing the time required to develop new cultivars. In addition plant  
5 genetic engineering permits the manufacture of products not naturally obtainable from flax, for example therapeutic agents.

In order to develop novel flax cultivars through plant genetic engineering, control over the expression of the introduced foreign or non-native gene is of critical importance. The desired expression  
10 characteristics for the non-native gene, such as the level of expression of the non-native gene, the particular plant tissue or organ in which the non-native gene is expressed, and the particular time in the growth cycle of the plant at which the non-native gene is expressed, will vary depending on the application for which the plant line is developed. For  
15 example, the modification of the seed oil composition may require low levels of seed-specific expression of an enzyme involved in fatty acid metabolism at an early stage in seed development (see for example US Patent 5,420,034). On the other hand expression of a pharmaceutical protein could preferably require high levels of leaf-specific expression  
20 upon harvesting of the plant leaves (see for example, US Patent 5,929,304).

In order to manipulate the expression characteristics of non-native genes numerous factors can be influenced. One factor is the choice of the transcriptional promoter used. A wide range of plant compatible promoters is currently available and some of the better  
25 documented promoters include constitutive promoters such as the 35-S CaMV promoter (Rothstein *et al.* (1987), Gene 53: 153-161) and the ubiquitin promoter (US Patent 5,614,399), tissue specific promoters such as seed-specific promoters, for example the phaseolin promoter (Sengupta-Gopalan *et al.*, (1985), PNAS USA 82: 3320-3324) and inducible  
30 promoters, such as those inducible by heat (Czarnencka *et al.*, (1989), Mol. Cell. Biol. 9 (8): 3457-3464), UV light, elicitors and wounding (Lois *et al.*, (1989) EMBO J. 8 (6): 1641-1648), or chemicals such as endogenous

005280" E654950

hormones (Skriver *et al.* (1991), Proc. Natl. Acad. Sci. USA 88(16): 7266-7270). Other factors that can be manipulated in order to control the expression characteristics of non-native gene in plants include transcriptional modification factors such as introns, polyadenylation sites and transcription termination sites. The expression characteristics of the non-native gene can further be manipulated by factors that affect translation, such as ribosomal binding sites and the codon bias that is exhibited by the host. Furthermore, the non-native gene itself may affect the viability of the transgenic plant, thus limiting particularly the levels of expression that can be attained. In some cases it may be possible to overcome this problem, by expressing the protein in a tissue specific manner, e.g. in the leaves or seed, or by restricting the accumulation of the protein in different subcellular compartments such as for example the cytoplasm, the endoplasmic reticulum or vacuoles, typically by the presence or the absence of specific targeting sequences capable of directing the protein to these compartments. Another factor that will affect the expression characteristics is the location in which the construct inserts itself into the host chromosome. This effect could provide an explanation as to why different plants, transformed with the same recombinant construct, can have fluctuating levels of recombinant protein expression.

To the best of the inventors' knowledge, expression of non-native genes in flax seeds is only documented in PCT Patent Application WO 98/18948. This application discloses two stearyl-acyl carrier protein desaturase (SAD) genes derived from flax. The associated SAD promoter sequences are useful for the modification of flax and other plants for the expression of endogenous or foreign genes. However the methods taught by WO 98/18948 are limited by the fact that the SAD promoters are not seed-specific in flax and confer expression to leaves, stems, flowers and seeds. Expression of non-native genes thus may result in undesirable side effects in non-seed tissues. In addition the use of the SAD promoters allows limited control over expression level and timing of expression.

There is a need in the art to further improve methods for the expression of non-native genes in flax seeds and other plant seeds.

#### SUMMARY OF THE INVENTION

The present invention relates to improved methods for the seed-specific expression of non-native genes in plants. In particular, the invention relates to improved methods for the seed-specific expression of non-native genes in flax.

Accordingly, in one aspect, the invention provides a method for the expression of a nucleic acid sequence of interest in flax seeds comprising:

- (a) preparing a chimeric nucleic acid construct comprising in the 5' to 3' direction of transcription as operably linked components
  - (1) a seed-specific promoter obtained from flax; and
  - (2) the nucleic acid sequence of interest wherein said nucleic acid of interest is non-native to said flax seed-specific promoter;
- (b) introducing said chimeric nucleic acid construct into a flax plant cell; and
- (c) growing said flax plant cell into a mature flax plant capable of setting seed, wherein said nucleic acid sequence of interest is expressed in the seed under the control of said flax seed-specific promoter.

In a preferred embodiment of the invention, at least one expression characteristic, e.g. timing of expression in the plant's life cycle, conferred by the promoter to the non-native nucleic acid sequence is similar to that expression characteristic when conferred to a native nucleic acid sequence. In further preferred embodiments, the flax seed-specific promoter is an oleosin promoter, a 2S storage protein promoter or a legumin-like seed storage protein promoter.

In a further aspect, the present invention provides transgenic flax seeds prepared according to a method comprising:

- (a) preparing a chimeric nucleic acid construct comprising in the 5' to 3' direction of transcription as operably linked components:

- (1) a seed-specific promoter obtained from flax; and
- (2) a nucleic acid sequence of interest wherein said nucleic acid of interest is non-native to said seed-specific promoter;

5 (b) introducing said chimeric nucleic acid construct into a flax plant cell; and

(c) growing said flax plant cell into a mature flax plant capable of setting seed, wherein said nucleic acid sequence of interest is expressed in the seed under the control of said seed-specific promoter.

10 In a further aspect the present invention provides flax plants capable of setting seed prepared by a method comprising:

(a) preparing a chimeric nucleic acid construct comprising in the 5' to 3' direction of transcription as operably linked components:

(1) a seed-specific promoter obtained from flax; and

15 (2) a nucleic acid sequence of interest wherein said nucleic acid of interest is non-native to said seed-specific promoter;

(b) introducing said chimeric nucleic acid construct into a flax plant cell; and

20 (c) growing said flax plant cell into a mature flax plant capable of setting seed, wherein said nucleic acid sequence of interest is expressed in the seed under the control of said seed-specific promoter.

In yet a further aspect, the present invention provides novel flax seed specific promoters useful for the expression of non-native genes  
25 in flax seeds and the seeds of other plant species, useful for example for modification of the protein or oil composition of the seed.

In a preferred embodiment, the seed specific promoter comprises:

(a) a nucleic acid sequence as shown in Figure 1  
30 (SEQ.ID.NO.:1), Figure 2 (SEQ.ID.NO.:4), Figure 3 (SEQ.ID.NO.:6) or Figure 4 (SEQ.ID.NO.:8) wherein T can also be U;

005280.E654960

(b) a nucleic acid sequence that is complimentary to a nucleic acid sequence of (a);

(c) a nucleic acid sequence that has substantial sequence homology to a nucleic acid sequence of (a) or (b);

5 (d) a nucleic acid sequence that is an analog of a nucleic acid sequence of (a), (b) or (c); or

(e) a nucleic acid sequence that hybridizes to a nucleic acid sequence of (a), (b), (c) or (d) under stringent hybridization conditions.

In another aspect, the invention provides chimeric nucleic acid  
10 sequences comprising a first nucleic acid sequence obtained from flax operatively linked to a second nucleic acid sequence non-native to said first nucleic acid sequence wherein said first nucleic acid sequence comprises a novel flax seed-specific promoter.

Other features and advantages of the present invention will  
15 become readily apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples while indicating preferred embodiments of the invention are given by way of illustration only, since various changes and modifications within the spirit and scope of the invention will become apparent to those  
20 skilled in the art of this detailed description.

#### **BRIEF DESCRIPTION OF THE DRAWINGS**

The invention will now be described in relation to the drawings in which:

Figure 1 shows the DNA sequence (SEQ.ID.NO.:1) of a flax  
25 genomic clone encoding a 16.0 kDa oleosin protein (SEQ.ID.NOS.:2 and 3).

Figure 2 shows the DNA sequence (SEQ.ID.NO.:4) of a flax genomic clone encoding a 18.6 kDa oleosin protein (SEQ.ID.NO.:5).

Figure 3 shows the DNA sequence (SEQ.ID.NO.:6) of a flax genomic clone encoding a 2S storage protein (SEQ.ID.NO.:7).

30 Figure 4 shows the DNA sequence (SEQ.ID.NO.:8) of a flax genomic clone encoding a 54.5 kDa legumin-like storage protein (SEQ.ID.NOS.:9-12).



Figure 5 shows Southern blot analysis of flax genomic DNA probed with flax oleosin DNA sequences.

Figure 6 shows a Northern blot analysis of the seed specific expression of flax oleosins.

5           Figure 7 shows a Northern blot analysis of the developmental expression of flax oleosins during seed development.

Figure 8 shows the GUS activity of flax embryos bombarded with flax oleosin promoter-GUS-flax terminator gene constructs.

10           Figure 9 shows GUS expression in developing flax embryos and Arabidopsis seeds of plants transformed with a 2S protein gene promoter GUS fusion.

Figure 10 shows the tissue-specific expression of GUS in transgenic flax plants transformed with a linin promoter-GUS-linin terminator gene construct.

15           Figure 11 shows the temporal expression of GUS in transgenic flax plants transformed a linin promoter-GUS-linin terminator gene construct.

20           Figure 12 shows the expression of GUS in transgenic *Brassica napus* plants (L1 to L9) transformed with a linin promoter-GUS-linin terminator gene construct.

Figure 13 shows the expression of GUS in transgenic Arabidopsis plants transformed with a linin promoter-GUS-linin terminator gene construct at different stages of seed development.

#### **DETAILED DESCRIPTION OF THE INVENTION**

25           As hereinbefore mentioned, the present invention relates to improved methods for the expression of non-native genes in plants, in particular flax. The invention provides methods allowing the seed-specific expression of non-native genes in flax. The methods of the invention are advantageous in that improved control over the expression  
30 of non-native genes in flax seeds is obtained. Expression of the non-native gene is restricted to the seeds, thereby limiting potential undesirable effects resulting from the expression in other plant organs or tissues. In addition,

the provided methodology allows improved control over expression characteristics, such as the expression level of the non-native gene and timing of expression of the non-native gene in the developmental cycle of the plant. The methods of the present invention are particularly useful in  
5 that in accordance with the present invention the seed composition with respect to valuable raw materials, such as oil, protein and polysaccharides, may be altered both qualitatively and quantitatively.

Accordingly, in one aspect, the invention provides a method for the expression of a nucleic acid sequence of interest in flax seeds  
10 comprising:

- (a) preparing a chimeric nucleic acid construct comprising in the 5' to 3' direction of transcription as operably linked components;
  - (1) a seed-specific promoter obtained from flax; and
  - (2) the nucleic acid sequence of interest wherein said nucleic  
15 acid of interest is non-native to said flax seed-specific promoter;
- (b) introducing said chimeric nucleic acid construct into a flax plant cell; and
- (c) growing said flax plant cell into a mature plant capable of  
20 setting seed, wherein said nucleic acid sequence of interest is expressed in the seed under the control of said flax seed-specific promoter.

As used herein, the term "non-native" refers to any nucleic acid sequence, including any RNA or DNA sequence, which is not normally associated with the seed-specific promoter. This includes  
25 heterologous nucleic acid sequences which are obtained from a different plant species as the promoter as well as homologous nucleic acid sequences which are obtained from the same plant species as the promoter but are not associated with the promoter in the wild-type (non-transgenic) plant.

30 The non-native nucleic acid sequence when linked to a seed-specific promoter obtained from flax results in a chimeric construct. The chimeric construct is introduced into a flax plant cell to create a

transgenic flax plant cell which results in a detectably different phenotype of the flax plant cell or flax plant grown from it when compared with a non-transgenic flax plant cell or flax plant grown from it. A contiguous nucleic acid sequence identical to the nucleic acid sequence of the chimeric construct is not present in the non-transformed flax plant cell or flax plant grown from it. In this respect, chimeric nucleic acid sequences include those sequences which contain a flax promoter linked to a nucleic acid sequence obtained from another plant species or a nucleic acid sequence from flax but normally not associated with that promoter. Chimeric nucleic acid sequences as used herein further include sequences comprising a flax promoter and a nucleic acid sequence that is normally linked to the promoter but additionally containing a non-native nucleic acid sequence. For example, if the promoter is a flax seed-specific oleosin promoter, sequences "non-native" to the flax oleosin promoter also include a sequence comprising a fusion between the flax oleosin gene naturally associated with the oleosin promoter, and a coding sequence of interest that is not naturally associated with the promoter. The term non-native is also meant to include a fusion gene as hereinabove which additionally includes a cleavage sequence separating the nucleic acid sequence that is normally linked to the promoter sequence and the gene encoding the protein of interest.

The term "nucleic acid sequence" refers to a sequence of nucleotide or nucleoside monomers consisting of naturally occurring bases, sugars and intersugar (backbone) linkages. The term also includes modified or substituted sequences comprising non-naturally occurring monomers or portions thereof, which function similarly. The nucleic acid sequences of the present invention may be ribonucleic (RNA) or deoxyribonucleic acids (DNA) and may contain naturally occurring bases including adenine, guanine, cytosine, thymidine and uracil. The sequences may also contain modified bases such as xanthine, hypoxanthine, 2-aminoadenine, 6-methyl, 2-propyl, and other alkyl adenines, 5-halo uracil, 5-halo cytosine, 6-aza uracil, 6-aza cytosine and

6-aza thymine, pseudo uracil, 4-thiouracil, 8-halo adenine, 8-amino adenine, 8-thiol adenine, 8-thio-alkyl adenines, 8-hydroxyl adenine and other 8-substituted adenines, 8-halo guanines, 8-amino guanine, 8-thiol guanine, 8-thioalkyl guanines, 8-hydroxyl guanine and other 8-substituted  
5 guanines, other aza and deaza uracils, thymidines, cytosines, adenines, or guanines, 5-trifluoromethyl uracil and 5-trifluoro cytosine.

The term "seed-specific promoter", means that a gene expressed under the control of the promoter is predominantly expressed in plant seeds with no or no substantial expression, typically less than 5% of  
10 the overall expression level, in other plant tissues.

In a further aspect, the present invention provides novel flax seed specific promoters useful for the expression of non-native genes in flax seeds and the seeds of other plant species. The promoters may be used to modify for example the protein, oil or polysaccharide composition of  
15 the seeds. In a preferred embodiment, the seed specific promoter comprises:

- (a) a nucleic acid sequence as shown in Figure 1 (SEQ.ID.NO.:1), Figure 2 (SEQ.ID.NO.:4), Figure 3 (SEQ.ID.NO.:6) or Figure 4 (SEQ.ID.NO.:8) wherein T can also be U;
- 20 (b) a nucleic acid sequence that is complimentary to a nucleic acid sequence of (a);
- (c) a nucleic acid sequence that has substantial sequence homology to a nucleic acid sequence of (a) or (b);
- (d) a nucleic acid sequence that is an analog of a nucleic acid  
25 sequence of (a), (b) or (c); or
- (e) a nucleic acid sequence that hybridizes to a nucleic acid sequence of (a), (b), (c) or (d) under stringent hybridization conditions.

The term "sequence that has substantial sequence homology" means those nucleic acid sequences which have slight or inconsequential sequence variations from the sequences in (a) or (b), i.e., the sequences  
30 function in substantially the same manner and are capable of driving seed specific expression of non-native nucleic acid sequences. The variations

may be attributable to local mutations or structural modifications. Nucleic acid sequences having substantial homology include nucleic acid sequences having at least 65%, more preferably at least 85%, and most preferably 90-95% identity with the nucleic acid sequences as shown in  
5 Figure 1 (SEQ.ID.NO.:1), Figure 2 (SEQ.ID.NO.:4), Figure 3 (SEQ.ID.NO.:6) or Figure 4 (SEQ.ID.NO.:8).

The term "sequence that hybridizes" means a nucleic acid sequence that can hybridize to a sequence of (a), (b), (c) or (d) under stringent hybridization conditions. Appropriate "stringent hybridization  
10 conditions" which promote DNA hybridization are known to those skilled in the art, or may be found in Current Protocols in Molecular Biology, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. For example, the following may be employed: 6.0 x sodium chloride/sodium citrate (SSC) at about 45°C, followed by a wash of 2.0 x SSC at 50°C. The stringency may be  
15 selected based on the conditions used in the wash step. For example, the salt concentration in the wash step can be selected from a high stringency of about 0.2 x SSC at 50°C. In addition, the temperature in the wash step can be at high stringency conditions, at about 65°C.

The term "a nucleic acid sequence which is an analog" means a  
20 nucleic acid sequence which has been modified as compared to the sequence of (a), (b) or (c) wherein the modification does not alter the utility of the sequence (i.e. as a seed specific promoter) as described herein. The modified sequence or analog may have improved properties over the sequence shown in (a), (b) or (c). One example of a modification to prepare  
25 an analog is to replace one of the naturally occurring bases (i.e. adenine, guanine, cytosine or thymidine) of the sequence shown in Figure 1, Figure 2, Figure 3 or Figure 4 with a modified base such as such as xanthine, hypoxanthine, 2-aminoadenine, 6-methyl, 2-propyl and other alkyl adenines, 5-halo uracil, 5-halo cytosine, 6-aza uracil, 6-aza cytosine and  
30 6-aza thymine, pseudo uracil, 4-thiouracil, 8-halo adenine, 8-aminoadenine, 8-thiol adenine, 8-thiolalkyl adenines, 8-hydroxyl adenine and other 8-substituted adenines, 8-halo guanines, 8 amino

guanine, 8-thiol guanine, 8-thiolalkyl guanines, 8-hydroxyl guanine and other 8-substituted guanines, other aza and deaza uracils, thymidines, cytosines, adenines, or guanines, 5-trifluoromethyl uracil and 5-trifluoro cytosine.

5 Another example of a modification is to include modified phosphorous or oxygen heteroatoms in the phosphate backbone, short chain alkyl or cycloalkyl intersugar linkages or short chain heteroatomic or heterocyclic intersugar linkages in the nucleic acid molecule shown in Figure 1, Figure 2, Figure 3 or Figure 4. For example, the nucleic acid  
10 sequences may contain phosphorothioates, phosphotriesters, methyl phosphonates, and phosphorodithioates.

A further example of an analog of a nucleic acid molecule of the invention is a peptide nucleic acid (PNA) wherein the deoxyribose (or ribose) phosphate backbone in the DNA (or RNA), is replaced with a  
15 polyamide backbone which is similar to that found in peptides (P.E. Nielsen, et al Science 1991, 254, 1497). PNA analogs have been shown to be resistant to degradation by enzymes and to have extended lives *in vivo* and *in vitro*. PNAs also bind stronger to a complimentary DNA sequence due to the lack of charge repulsion between the PNA strand and the DNA  
20 strand. Other nucleic acid analogs may contain nucleotides containing polymer backbones, cyclic backbones, or acyclic backbones. For example, the nucleotides may have morpholino backbone structures (U.S. Pat. No. 5,034,506). The analogs may also contain groups such as reporter groups, a group for improving the pharmacokinetic or pharmacodynamic properties  
25 of nucleic acid sequence.

In another aspect, the invention provides chimeric nucleic acid sequences comprising a first nucleic acid sequence obtained from flax operatively linked to a second nucleic acid sequence non-native to said first nucleic acid sequence wherein said first nucleic acid sequence  
30 comprises a novel flax seed-specific promoter. Preferably, the promoter is selected from the group of promoters comprising Figure 1, Figure 2, Figure

3 and Figure 4 or a nucleic acid sequence hybridizing thereto under stringent conditions.

In accordance with the present invention, the chimeric nucleic acid sequences can be incorporated in a known manner in a recombinant expression vector which ensures good expression in the seed cell. Accordingly, the present invention includes a recombinant expression vector comprising a chimeric nucleic acid sequence of the present invention suitable for expression in a seed cell.

The term "suitable for expression in a seed cell" means that the recombinant expression vectors contain the chimeric nucleic acids sequence of the invention, a regulatory region and a termination region, selected on the basis of the seed cell to be used for expression, which is operatively linked to the nucleic acid sequence encoding the polypeptide of desirable amino acid composition. Operatively linked is intended to mean that the chimeric nucleic acid sequence encoding the polypeptide is linked to a regulatory sequence and termination region which allows expression in the seed cell. A typical construct consists, in the 5' to 3' direction of a regulatory region complete with a promoter capable of directing expression in a plant, a polypeptide coding region and a transcription termination region functional in plant cells. These constructs may be prepared in accordance with methodology well known to those of skill in the art of molecular biology (see for example: Sambrook *et al.* (1990), Molecular Cloning, 2<sup>nd</sup> ed. Cold Spring Harbor Press). The preparation of constructs may involve techniques such as restriction digestion, ligation, gel electrophoresis, DNA sequencing and PCR. A wide variety of cloning vectors is available to perform the necessary cloning steps. Especially suitable for this purpose are the cloning vectors with a replication system that is functional in *Escherichia coli* such as pBR322, the pUC series M13mp series, pACYC184, pBluescript etc. Nucleic acid sequences may be introduced into these vectors and the vectors may be used to transform *E. coli* which may be grown in an appropriate medium. Plasmids may be recovered from the cells upon harvesting and lysing the cells. Final

constructs may be introduced into plant vectors compatible with integration into the plant such as the Ti and Ri plasmids.

The methods for the expression of non-native genes in flax seeds in accordance with the present invention may be practiced using any  
5 flax seed-specific promoter and are not limited by the specific flax seed specific promoter that is selected. In preferred embodiments of the present invention, the flax seed-specific promoter confers to the non-native nucleic acid sequence at least one expression characteristic which is similar or identical to an expression characteristic conferred to the native nucleic  
10 acid sequence by the native promoter. The term "expression characteristic" as used herein refers to any measurable property or effect conferred by the flax seed-specific promoter to the nucleic acid sequence operably linked to the flax seed-specific promoter. Thus in preferred embodiments, timing of expression in the plant's life cycle, of the  
15 non-native nucleic acid sequence is similar or identical to timing of expression of the native nucleic acid sequence. In further preferred embodiments, the expression level of the heterologous nucleic acid sequence is similar or identical to the expression level of the native nucleic acid sequence. In yet further specific embodiments, the response of  
20 the non-native gene to alterations in lighting conditions, changes in wavelength or light intensity for example, changes in temperature, tissue wounding, changes in concentration of chemical agents, such as for example phytohormones and pesticides, is similar to the response of the native nucleic acid sequence to these stimuli. Other desired expression  
25 characteristics conferred by a flax seed-specific promoter may be recognized by those skilled in the art and a flax seed-specific promoter may be selected accordingly.

Flax-seed specific promoters that may be used in accordance with the present invention include promoters associated with seed storage  
30 proteins, such as all albumins and globulins, including the vicilin and legumin-like proteins, as well as seed-specific promoters not associated with seed storage proteins, such as oleosins. Of further particular interest



are promoters associated with fatty acid metabolism, such as acyl carrier protein (ACP), saturases, desaturases, elongases and the like.

In preferred embodiments of the present invention the seed specific promoter used is an oleosin promoter, a legumin-like seed storage protein promoter or a 2S storage protein promoter. In particularly preferred embodiments the seed specific promoter has the sequence shown in Figure 1, Figure 2, Figure 3 or Figure 4 or any nucleic acid sequences obtainable from flax and hybridizing to any one of these four nucleic acid sequences under stringent conditions.

Additional flax seed-specific promoters may be used in accordance with the present invention. These promoters may be obtained in a number of ways. Where a flax seed protein has been isolated, it may be partially sequenced, so that a nucleic acid probe may be designed for identifying RNA specific to the seed. To further enhance the RNA specifically associated with the seed, cDNA may be prepared from seed cells and the cDNA may be subtracted with mRNA or cDNA from non-seed cells. The remaining seed cDNA may then be used to probe a genomic DNA library for complementary sequences. Sequences hybridizing to the cDNA may subsequently be obtained and the associated promoter region may be isolated. It is also possible to screen genomic DNA libraries prepared from flax seed tissues using known seed specific genes from other plant species and subsequently isolate their associated promoters. Due to the relative abundance of seed-storage proteins in seeds it is also be possible to obtain sequence information through random sequencing of flax seed cDNA libraries. Those cDNA sequences matching sequence of known seed-storage proteins could be used to identify the associated promoter. Databases containing sequence information from large scale sequencing from for example *Arabidopsis* and maize may be searched for known seed-specific proteins and/or promoters and the information may be used to identify promoter sequences in flax that share sequence similarity. Alternative methods to isolate additional flax seed specific promoters may be used and novel flax seed specific promoters may be

discovered by those skilled in the art and used in accordance with the present invention.

5 The nucleic acid sequence of interest linked to the promoter may be any nucleic acid sequence of interest including any RNA or DNA sequence encoding a peptide or protein of interest, for example, an enzyme, or a sequence complementary to a genomic sequence, where the genomic sequence may be at least one of an open reading frame, an intron, a non-coding leader sequence, or any sequence where the complementary sequence will inhibit transcription, messenger RNA processing, for  
10 example splicing or translation. The nucleic acid sequence of interest may be synthetic, naturally derived or a combination thereof. As well, the nucleic acid sequence of interest could be a fragment of the natural sequence, for example just include the catalytic domain or a structure of particular importance. Depending upon the nature of the nucleic acid  
15 sequence of interest, it may be desirable to synthesize the sequence with plant preferred codons. The plant preferred codons may be determined from the codons of highest frequency in the proteins expressed in the largest amount in particular plant species of interest.

20 The nucleic acid sequence of interest may encode any of a variety of recombinant proteins. Examples of recombinant proteins which might be expressed by the methods of the present invention include proteins with a favorable catalytic function or a valuable protein that will accumulate to high levels and be extracted if desired. Proteins with a catalytic function, include, but are not limited to, proteins that confer a  
25 new biochemical phenotype on the developing seeds. New phenotypes could include such modifications as altered seed-protein or seed oil composition or seed polysaccharide composition, enhanced production of pre-existing desirable products or properties and the reduction or even suppression on an undesirable gene product using antisense, ribozyme or  
30 co-suppression technologies (Izant and Weintraub (1984) Cell 26: 1007-1015, antisense; Hazelhoff and Gerlach (1988) Nature 334: 585-591, ribozyme; Napoli *et al.* (1990) Plant Cell 2: 279-289, co-suppression).

It is expected that the desired proteins would be expressed in all embryonic tissues, although varying cellular expression may be detected in the different embryonic tissues such as the embryonic axis and cotyledons. The nucleic acid sequence of interest may be expressed at any stage in seed  
5 development. The timing of expression may depend on the particular use of the invention. Expression of enzymes involved in oil modification may be desirable early in seed development, for example before accumulation of seed storage protein.

Besides the promoter region and the nucleic acid sequence of  
10 interest, a nucleic acid sequence capable of terminating transcription is typically included in expression vectors. Transcriptional terminators are preferably about 200 to about 1,000 nucleotide base pairs and may comprise any such sequences functional in plants, such as the nopaline synthase termination region (Bevan *et al.*, (1983) Nucl. Acid. Res. 11: 369-385), the  
15 phaseolin terminator (van der Geest *et al.*, (1994) Plant J. 6(3): 413-423), the terminator for the octopine synthase gene of *Agrobacterium tumefaciens* or other similarly functioning elements. These transcription terminator regions can be obtained as described by An (1987), Methods in Enzym. 153: 292 or are already present in plasmids available from commercial sources  
20 such as ClonTech, Palo Alto, California. The choice of the appropriate terminator may have an effect of the rate of transcription.

The chimeric construct may further comprise enhancers such as the AMV leader (Jobling and Gehrke (1987), Nature 325: 622-625) or introns. It should be understood that the design of the expression vector  
25 may depend on such factors as the choice of the plant species and/or the type of polypeptide to be expressed.

The expression vectors will normally also contain a marker gene. Marker genes comprise all genes that enable distinction of transformed plant cells from non-transformed cells, including selectable  
30 and screenable marker genes. Conveniently, a marker may be a resistance marker to a herbicide, for example, glyphosate or phosphinothricin, or to an antibiotic such as kanamycin, G418, bleomycin, hygromycin,

chloramphenicol and the like, which confer a trait that can be selected for by chemical means. Screenable markers may be employed to identify transformants through observation. They include but are not limited to the  $\beta$ -glucuronidase or *uidA* gene, a  $\beta$ -lactamase gene or a green fluorescent protein (Niedz *et al.* (1995) Plant Cell Rep. 14: 403).

In order to introduce nucleic acid sequences into plant cells in general a variety of techniques are available to the skilled artisan. *Agrobacterium*-mediated transformation for flax plant cells has been reported and flax transformants may be obtained in accordance with the methods taught by Dong and McHughen (1993) Plant Science 88: 61-77, although a variety of other techniques (see below) may also be used to introduce the chimeric DNA constructs in flax cells if so desired.

Transformed flax plants grown in accordance with conventional agricultural practices known to a person skilled in the art are allowed to set seed. Flax seed may then be obtained from mature flax plants and analyzed for desired altered properties with respect to the wild-type seed.

Two or more generations of plants may be grown and either crossed or selfed to allow identification of plants and strains with desired phenotypic characteristics including production of the recombinant polypeptide. It may be desirable to ensure homozygosity in the plants to assure continued inheritance of the recombinant trait. Methods for selecting homozygous plants are well known to those skilled in the art of plant breeding and include recurrent selfing and selection and anther and microspore culture. Homozygous plants may also be obtained by transformation of haploid cells or tissues followed by regeneration of haploid plantlets subsequently converted to diploid plants by any number of known means (e.g. treatment with colchicine or other microtubule disrupting agents).

The present invention also includes transgenic flax seeds prepared according to a method comprising:

(a) preparing a chimeric nucleic acid construct comprising in the 5' to 3' direction of transcription as operably linked components:

- (1) a seed-specific promoter obtained from flax; and
- (2) a nucleic acid sequence of interest wherein said nucleic acid of interest is non-native to said seed-specific promoter;

(b) introducing said chimeric nucleic acid construct into a flax plant cell; and

(c) growing said flax plant cell into a mature flax plant capable of setting seed

wherein said nucleic acid sequence of interest is expressed in the seed under the control of said seed-specific promoter.

In preferred embodiments of the invention the seed-specific promoter is selected from the group of flax seed specific promoters consisting of, a 2S storage protein promoter, a globulin promoter, an oleosin promoter, and a legumin-like seed storage protein promoter. Specific promoter sequences are shown in Figure 1 (SEQ.ID.NO.:1), Figure 2 (SEQ.ID.NO.:4), Figure 3 (SEQ.ID.NO.:6) and Figure 4 (SEQ.ID.NO.:8).

The present invention further provides flax plants capable of setting seed prepared by a method comprising:

(a) preparing a chimeric nucleic acid construct comprising in the 5' to 3' direction of transcription as operably linked components:

- (1) a seed-specific promoter obtained from flax; and
- (2) a nucleic acid sequence of interest wherein said nucleic acid of interest is non-native to said seed-specific promoter;

(b) introducing said chimeric nucleic acid construct into a flax plant cell; and

(c) growing said flax plant cell into a mature flax plant capable of setting seed

wherein said nucleic acid sequence of interest is expressed in the seed under the control of said seed-specific promoter.

The present invention further provides methods of use for the novel promoters shown in Figure 1 (SEQ.ID.NO.:1), Figure 2 (SEQ.ID.NO.:4), Figure 3 (SEQ.ID.NO.:6) and Figure 4 (SEQ.ID.NO.:8) in plant species other than flax. Accordingly, the invention also includes the preparation of chimeric nucleic acid constructs comprising a promoter selected from the group promoters shown in Figure 1, Figure 2, Figure 3 and Figure 4 and a nucleic acid sequence of interest, and expression in a seed-specific manner of the nucleic acid sequence of interest in plant species other than flax and under the control of the flax promoter.

In another aspect of the present invention there is provided a method for the expression of a nucleic acid sequence of interest in plant seeds comprising:

- (a) preparing a chimeric nucleic acid construct comprising in the 5' to 3' direction of transcription as operably linked components;
  - (1) a seed-specific promoter selected from the group of seed-specific promoters consisting of
    - (i) a nucleic acid sequence as shown in Figure 1 (SEQ.ID.NO.:1), Figure 2 (SEQ.ID.NO.:4), Figure 3 (SEQ.ID.NO.:6) or Figure 4 (SEQ.ID.NO.:8) wherein T can also be U;
    - (ii) a nucleic acid sequence that is complimentary to a nucleic acid sequence of (i);
    - (iii) a nucleic acid sequence that has substantial sequence homology to a nucleic acid sequence of (i) or (ii); and
    - (iv) a nucleic acid sequence that is an analog of a nucleic acid sequence of (i), (ii) or (iii);
    - (v) a nucleic acid sequence that hybridizes to a nucleic acid sequence of (i), (ii), (iii) or (iv) under stringent hybridization conditions; and
  - (2) said nucleic acid of interest;
  - (b) introducing the chimeric nucleic acid construct into a plant cell;

(c) growing said plant cell into a mature plant capable of setting seed, wherein said nucleic acid sequence of interest is expressed in the seed under the control of said seed-specific promoter.

5 A variety of techniques are available for the introduction of nucleic acid sequences, in particular DNA, into plant host cells in general. For example, the chimeric DNA constructs may be introduced into host cells obtained from dicotyledonous plants, such as tobacco, and oleoagenous species, such as *Brassica napus* using standard *Agrobacterium* vectors by a transformation protocol such as described by Moloney *et al.* 10 (1989), Plant Cell Rep. 8: 238-242 or Hinchey *et al.* (1988) Bio/Technol. 6: 915-922; or other techniques known to those skilled in the art. For example, the use of T-DNA for transformation of plant cells has received extensive study and is amply described in EP 0 120 516, Hoekema *et al.*, (1985), Chapter V In: *The Binary Plant Vector System* Offset-drukkerij 15 Kanters BV, Alblaserdam); Knauf *et al.* (1983), *Genetic Analysis of Host Expression by Agrobacterium*, p. 245, In: *Molecular Genetics of Bacteria-Plant Interaction*, Puhler, A. ed. Springer-Verlag, NY); and An *et al.*, (1985), (EMBO J., 4: 277-284). *Agrobacterium* transformation may also be used to transform monocot plant species (US Patent 5,591,616).

20 Conveniently, explants may be cultivated with *Agrobacterium tumefaciens* or *Agrobacterium rhizogenes* to allow for the transfer of the transcription construct in the plant host cell. Following transformation using *Agrobacterium* the plant cells are dispersed into an appropriate medium for selection, subsequently callus, shoots and eventually plants 25 are recovered. The *Agrobacterium* host will harbour a plasmid comprising the vir genes necessary for transfer of the T-DNA to plant cells. For injection and electroporation (see below) disarmed Ti-plasmids (lacking the tumour genes, particularly the T-DNA region) may be introduced into the plant cell.

30 The use of non-*Agrobacterium* techniques permits the use of constructs described herein to obtain transformation and expression in a wide variety of monocotyledonous and dicotyledonous plant species.

These techniques are especially useful for transformation of plant species that are intractable in an *Agrobacterium* transformation system. Other techniques for gene transfer include particle bombardment (Sanford, (1988), Trends in Biotechn. 6: 299-302), electroporation (Fromm *et al.*, (1985), PNAS USA, 82: 5824-5828; Riggs and Bates, (1986), PNAS USA 83: 5602-5606), PEG mediated DNA uptake (Potrykus *et al.*, (1985), Mol. Gen. Genetics., 199: 169-177), microinjection (Reich *et al.*, Bio/Techn. (1986) 4:1001-1004) and silicone carbide whiskers (Kaeppler *et al.* (1990) Plant Cell Rep. 9: 415-418).

10 In a further specific applications such as to *B. napus*, the host cells targeted to receive recombinant DNA constructs typically will be derived from cotyledonary petioles as described by Moloney *et al.* (1989) Plant Cell Rep. 8: 238-242. Other examples using commercial oil seeds include cotyledon transformation in soybean explants (Hinchee *et al.*, 15 (1988) Bio/Technol. 6: 915-922) and stem transformation of cotton (Umbeck *et al.*, (1987) Bio/Technol. 5: 263-266).

Following transformation, the cells, for example as leaf discs, are grown in selective medium. Once the shoots begin to emerge, they are excised and placed onto rooting medium. After sufficient roots have 20 formed, the plants are transferred to soil. Putative transformed plants are then tested for presence of a marker. Southern blotting is performed on genomic DNA using an appropriate probe, to show integration into the genome of the host cell.

The methods provided by the present invention can be used in 25 conjunction a broad range of plant species. Particularly preferred plant cells employed in accordance with the present invention include cells from the following plants: soybean (*Glycine max*), rapeseed (*Brassica napus*, *Brassica campestris*), sunflower (*Helianthus annuus*), cotton (*Gossypium hirsutum*), corn (*Zea mays*), tobacco (*Nicotiana tabacum*), 30 alfalfa (*Medicago sativa*), wheat (*Triticum* sp.), barley (*Hordeum vulgare*), oats (*Avena sativa* L.), sorghum (*Sorghum bicolor*), *Arabidopsis thaliana*, potato (*Solanum* sp.), flax/linseed (*Linum usitatissimum*), safflower



(*Carthamus tinctorius*), oil palm (*Eleais guineensis*), groundnut (*Arachis hypogaea*), Brazil nut (*Bertholletia excelsa*) coconut (*Cocos nucifera*), castor (*Ricinus communis*), coriander (*Coriandrum sativum*), squash (*Cucurbita maxima*), jojoba (*Simmondsia chinensis*) and rice (*Oryza sativa*).

5           The present invention has a variety of uses which include improving the intrinsic value of plant seeds by their accumulation of altered polypeptides or novel recombinant peptides or by the incorporation or elimination or a metabolic step. Use of the invention may result in improved protein quality (for example, increased  
10 concentrations or essential or rare amino acids), improved liquid quality by a modification of fatty acid composition, or improved or elevated carbohydrate composition. Examples include the expression of sulfur-rich proteins, such as those found in lupins or brazil nuts in a seed deficient in sulphurous amino acids. Improved protein quality could also be achieved  
15 by the expression of a protein or a fragment of a protein that is enriched in essential amino acids including lysine, cysteine, methionine and tryptophan. Alternatively, a fatty acyl coenzyme A, a transferase enzyme capable of modifying fatty acid ratios in triglycerides (storage lipid) could be expressed. In cases where a recombinant protein is allowed to accumulate  
20 in the seed, the protein could also be a peptide which has pharmaceutical or industrial value. In this case the peptide could be extracted from the seed and used in crude or purified form as appropriate for the intended use. As well, the polypeptides that are expressed in the seeds can be fragments or derivatives or the native protein. Pharmaceutically useful  
25 proteins may include, but are not limited to, anticoagulants, such as hirudin, antibodies, including monoclonal antibodies and antibody fragments, vaccines, cytokines or growth factors such as bovine growth factor, cholinergic differentiation factor (CDF), ciliary neurotrophic factor (CNTF), fibroblast growth factor (FGF), fish growth factor, gonadotropin,  
30 granulocyte-colony stimulating factor (G-CSF), granulocyte-macrophage colony-stimulating factor (GM-CSF), human growth hormone, interferon alpha (IFN- $\alpha$ ), interferon beta (IFN- $\beta$ ), interferon gamma (IFN- $\gamma$ ),

interleukin 1-alpha (IL1- $\alpha$ ), interleukin 1-beta (IL1- $\beta$ ), interleukin-2 (IL-2), interleukin-3 (IL-3), interleukin-4 (IL-4), interleukin-5 (IL-5), interleukin-6 (IL-6), interleukin-10 (IL-10), leukemia inhibitory factor (LIF), thioredoxin, macrophage colony-stimulating factor (M-CSF), myelomonocytic growth factor, nerve growth factor (NGF), oncostatin M, platelet-derived growth factor (PDGF), prolactin, transforming growth factor alpha (TGF- $\alpha$ ), transforming growth factor beta2 (TGF- $\beta$ 2), tumour necrosis factor alpha (TNF- $\alpha$ ), and tumour necrosis factor beta (TNF- $\beta$ ). Pharmaceutically useful proteins can also include mammalian proteins, for example, but not limited to  $\alpha$ -1-antitrypsin, anti-obesity proteins, blood proteins, collagen, collagenase, elastin, elastase, enteropeptidase, fibrinogen, haemoglobin, human serum albumin, insulin, lactoferrin, myoglobin and pulmonary surfactant proteins.

Industrially useful peptides may include, but are not limited to  $\alpha$ -amylase or other amylases, amyloglucosidase, arabinase, catalase, cellobiohydrolase, cellulases, chitinases, chymotrypsin, dehydrogenases, endo-glucanase, chymosin, endo-galactanase, esterases,  $\beta$ -galactosidase,  $\alpha$ -galactosidase or other galactosidases, gastric lipases, glucanases, glucose isomerase, hemi-cellulases, hydrolases, isomerase, ligninases, lipases, lyases, lysozymes, oxidases, oxidoreductase, papain, pectinases, pectin lyase, peroxidases, phosphatases, phytase, proteases, pullulanases, reductases, serine proteases, thioredoxin, transferase, trypsin, and xylanase.

The following non-limiting examples are illustrative of the present invention:

## **EXAMPLES**

### **EXAMPLE 1**

#### **Isolation of Seed-Specific Flax Promoters**

Seed specific cDNA clones were isolated from a flax seed specific cDNA-library. These cDNA clones were sequenced and the Basic Local Alignment Search Tool (BLAST) was used to compare these sequences against others in public databases such as Genbank. This

comparison revealed that the deduced amino acid sequence of several of the isolated cDNAs had a high degree of similarity to both the low and high molecular weight class of oleosins, 2S-albumin and legumin-like storage proteins. Probes were prepared individually from (portions of) cDNAs encoding oleosins, 2S albumin and legumin-like storage proteins and these were used to screen a genomic library prepared from the flax line Forge that is homozygous for four rust resistance genes (Anderson *et al.* (1997), The Plant Cell 9: 641-651). Several positive lambda clones for each probe were identified after high-stringency screening. The inserts were subcloned into the plasmid vector pBluescript and sequenced. Sequence information revealed that we had isolated the genomic counterparts to the oleosins, 2S albumin and cDNAs *legumin-like* cDNAs. Sequence information of the genomic clones containing sequences encoding a high and low molecular weight oleosin isoforms, 2S albumin and a legumin-like gene are presented in Figures 1 to 4 respectively.

Figure 1 and SEQ.ID.NO.:1 shows the DNA sequence of a flax genomic clone encoding a 16.0 kDa oleosin protein (low molecular weight or L-isoform). Putative regulatory elements are identified and indicated. These include inverted repeats (base pairs 805 to 813 and 821 to 829; base pairs 1858 to 1866 and 1877 to 1885), direct repeats (base pairs 184 to 193 and 1102 and 1111; base pairs 393 to 402 and 1701 to 1710; base pairs 683 to 692 and 1546 to 1555; base pairs 770 to 781 and 799 to 810; base pairs 955 to 964 and 1936 to 1945; base pairs 1483 to 1496 and 1513 to 1526), the abscisic acid responsive element (ABRE) (base pairs 1859 to 1866), CACA box (base pairs 1933 to 1936), TATA box (base pairs 1925 to 1931) and CAT box (base pairs 1989 to 1993). As well, the poly adenylation signal is indicated (base pairs 3020 to 3025). The open reading frame is interrupted by 1 short intron (which are marked) and the 2 exons are translated and indicated in IUPAC single letter amino-acid codes.

Figure 2 and SEQ.ID.NO.:4 shows the DNA sequence of a flax genomic clone encoding a 18.6 kDa oleosin protein (high molecular weight or H-isoform). Putative regulatory elements are identified and indicated.

These include direct repeats (base pairs 14 to 25 and 1427 to 1438; base pairs 80 to 89 and 1242 to 1251; base pairs 177 to 186 and 837 to 846; base pairs 1281 to 1290 and 1242 to 1251; base pairs 1591 to 1600 and 1678 to 1287). The open reading frame is not interrupted by introns and is translated and indicated in IUPAC single letter amino-acid codes.

Figure 3 and SEQ.ID.NO.:6 shows the DNA sequence of the flax genomic clone encoding a 2S storage protein. Nucleotide sequencing of this clone revealed it to have an open reading frame of 174 amino acids that showed homology to the plant 2S storage group of proteins. The sequence encodes an open reading frame with 38% overall similarity to a *Brassica oleracea* 2S storage protein, including complete conservation of the glutamine-rich stretch QQQGQQQGQQQ (SEQ.ID.NO.:13). In addition, the 2S storage protein gene promoter contained several putative promoter regulatory elements. These include AT rich repeats (base pairs 25-36, 97-108 and 167-190), RY-like repeat (base pairs 240-247), G-box-like element (base pairs 274-280), seed specific box-like motif (base pairs 285-290) and TATA box (base pairs 327-333).

Figure 4 and SEQ.ID.NO.:8 shows the DNA sequence of a flax genomic clone encoding a 54.4 kDa flax legumin-like seed storage protein. This legumin-like seed storage protein gene will also be referred to as "linin". The deduced amino acid sequence of the linin gene was compared to the *legumin-like* protein from *R. communis*, the legumin precursor from *M. salicifolia*, *Q. robur* and *G. hirsutum*, the glutelin precursor from *O. sativa* and a 12 S seed storage protein from *A. thaliana*. The linin gene shows a sequence identity/similarity with the corresponding proteins from *R. communis*, *M. salicifolia*, *Q. robur*, *G. hirsutum*, *O. sativa* and *A. thaliana* of 59/15, 47/16, 50/17, 45/17, 43/18 and 43/18 percent respectively. Putative regulatory elements in the promoter region are identified and indicated. These include inverted repeats (base pairs 265 to 276 and 281 to 292; base pairs 513 to 524 and 535 to 545), repeats (base pairs 1349 to 1360 and 1367 to 1378; base pairs 1513 to 1529 and 1554 to 1572), the abscisic acid responsive element (ABRE) (base pairs 1223 and 1231), legumin box (RY

repeats) (between base pairs 1223 and 1231), a possible vicilin box region (base pairs 1887 to 1894), CAAT box (base pairs 1782 to 1785) and TATA box (base pairs 1966 to 1970). As well, the signal peptide for ER membrane targeting is indicated (base pairs 2034-2080). The open reading frame is interrupted by 3 short introns (which are marked) and the 4 exons are translated and indicated in IUPAC single letter amino-acid codes.

Figure 5 shows Southern blot analysis of flax genomic DNA. 60 µg of flax genomic DNA was isolated from leaves, digested with EcoRI (lane 1), HindIII (lane 2) and BamHI (lane 3) and was loaded into the respective lanes. A) Hybridizations were performed with random primed <sup>32</sup>P-labelled 3T cDNA (high molecular weight flax oleosin isoform). B) Hybridizations were performed with random primed <sup>32</sup>P-labelled 7R cDNA (low molecular weight flax oleosin isoform). The results demonstrate that both 3T (high molecular weight oleosin isoform) and 7R (low molecular weight oleosin isoform) oleosin cDNAs hybridize with flax genomic DNA. More specifically the results indicate that 3T is likely to represent a 2-copy gene in flax, as seen by two bands in each lane of digestion. Similarly, 7R is likely to represent a multigene family in flax as multiple bands were detected for each digestion.

## 20 **EXAMPLE 2**

### **Seed specific expression of flax oleosin genes**

Figure 6 shows a Northern blot analysis of the seed specific expression of flax oleosins. Northern hybridization of the two oleosin mRNA in different tissues. Ten µg of total RNA was extracted from different tissues, R, root; C, cotyledon; L, leaf; S, seed capsule; E, embryo. The membrane was probed with (A) cDNA encoding high molecular weight (H)-isoform (identical to coding sequence as presented in Figure 2) and (B) cDNA encoding low molecular weight (L) -isoform (identical to coding sequence as presented in Figure 1). Both the transcripts are expressed only in the embryo and seed capsule, which contains embryos.

### **EXAMPLE 3**

#### **Developmental expression of flax oleosin genes during seed development**

Figure 7 shows a Northern blot analysis of the developmental expression of flax oleosins during seed development. 15 µg per lane of total RNA was loaded in each lane on agarose/formaldehyde gel and blotted onto HybondN+ membrane. 10J: This membrane was probed using the <sup>32</sup>P dCTP labeled flax oleosin cDNA clone (low molecular weight isoform). Stages indicated are the number of days past anthesis (DPA). 3T: 15 µg per lane of total RNA was loaded in each lane on agarose/formaldehyde gel and blotted onto HybondN+ membrane. 3T: This membrane was probed using the <sup>32</sup>P dCTP labeled flax oleosin cDNA clone (high molecular weight isoform). Both the transcripts were expressed very early in development (6DPA, early cotyledonary stage). Expression is maximum at 16 to 20 DPA (late cotyledonary stage) and declines at 22 DPA (mature embryos).

### **EXAMPLE 4**

#### **Transient Seed specific expression of β-glucuronidase (GUS) when under the regulatory control of flax oleosin regulatory sequences**

Two constructs were made using standard molecular biology techniques (eg see Sambrook *et al.* (1990), Molecular Cloning, 2nd ed. Cold Spring Harbor Press, including restriction enzyme digestions, ligation and polymerase chain reaction (PCR).

**Construct pSC54:** The β-glucuronidase reporter coding sequence from vector GUSN358>S (Clontech Laboratories) was placed between the promoter sequence from nucleotide 21 to 1852 and terminator sequence from 2395 to 3501 (as described in Figure 1). This insert was cloned into pBluescript and the resulting vector is called pSC54

**Construct pSC60:** The β-glucuronidase reporter coding sequence from vector GUSN358>S (Clontech Laboratories) was placed between the promoter sequence from nucleotide 1 to 2023 and terminator sequence from 2867 to 3925 (as described in Figure 2). This insert was cloned into pBluescript and the resulting vector is called pSC60.

pSC54, pSC60 and a promoter-less GUS construct (Control) were introduced into the flax embryos using particle bombardment using standard protocols (eg see Abenes *et al.* (1997) Plant Cell reports 17:1-7). Figure 8 shows the GUS activity of flax embryos bombarded with pSC54, pSC60 and a promoterless GUS construct measured 48 hours after particle bombardment. As can be seen the flax oleosin regulatory sequences are sufficient to drive the expression of GUS in flax embryos.

#### **EXAMPLE 5**

##### **Stable seed specific expression of $\beta$ -glucuronidase (GUS) in flax and *Arabidopsis* when under the regulatory control of flax 2S storage protein gene promoter**

A GUS reporter gene construct was made by incorporating 5' and 3' regions from the DNA fragment described in Figure 3 into promoterless-GUS pBI101 vector as follows.

A 400bp amplicon from the 5' end of the DNA fragment described in Figure 3 was PCR amplified using the following primers (location shown in Fig 3):

5' primer(1): 5'-TCCACTATGTAGGTCATA-3' (SEQ.ID.NO.:14)

3' primer(1): 5'-CTTTAAGGTGTGAGAGTC-3' (SEQ.ID.NO.:15)

The PCR primers also contained restriction sites for HindIII and BamHI which were used to clone the 400bp 5'UTR amplicon into the HindIII/BamHI sites of the pBI101 vector in front of the GUS reporter gene. A 736bp amplicon from the 3' untranslated region (3'UTR) of the DNA fragment described in Figure 3 was PCR amplified using the following primers (location shown in Fig 3):

5' primer (2):5'-AGGGGTGATCGATTA-3' (SEQ.ID.NO.:16)

3' primer (2):5'-GATAGAACCCACACGAGC-3' (SEQ.ID.NO.:17)

The PCR primers also contained restriction sites for SacI and EcoRI. The NOS terminator region of the pBI101 vector was cut out with SacI/EcoRI digestion and replaced with the similarly digested 736bp 3'UTR amplicon of the DNA fragment described in Figure 3.

The GUS reporter construct was then electroporated into *Agrobacterium tumifaciens* strain AGL1 and transformation of flax (Finnegan *et al.* (1993) Plant Mol Biol. 22(4): 625-633) and *Arabidopsis* (Valvekens *et al.* Proc. Natl. Acad. Sci. 85: 5536-5540) carried out according to previously described protocols.

Various tissues from flax and *Arabidopsis* plants carrying the GUS reporter construct were assayed histologically for evidence of GUS activity. In the case of flax, leaf tissue, root tissue and mid-maturity embryos dissected out of developing seeds were stained for GUS activity. For *Arabidopsis*, developing seeds were stained for GUS *in situ* in their siliques.

GUS staining was carried out by immersing the tissues in histochemical buffer containing 0.5 mM X-gluc, 0.5 M potassium phosphate buffer (pH 7.0), 1 mM EDTA, 0.5 M sorbitol, 0.5 mM potassium ferricyanide and 0.5 mM potassium ferrocyanide. The staining reaction was carried out for 12-16 hrs at 37°C and the reaction was stopped by adding 95% ethanol. Tissues were subsequently cleared of chlorophyll by repeated washing in 95% ethanol prior to photography. Figure 9 shows clear evidence of strong GUS activity in developing flax embryos and *Arabidopsis* seeds, and no evidence of GUS reporter gene expression in flax roots or leaves, or in *Arabidopsis* silique walls.

#### **EXAMPLE 6**

**Stable seed specific expression of  $\beta$ -glucuronidase (GUS) in flax, *Arabidopsis* and *Brassica napus* when under the regulatory control of flax legumin-like storage protein gene regulatory sequences**

A construct was made using standard molecular biology techniques, including restriction enzyme digestions, ligation and polymerase chain reaction (PCR). In order to obtain a DNA fragment containing approximately 2 kilobases from the 5' transcriptional initiation region of the flax legumin-like seed storage protein in a configuration suitable for ligation to a GUS coding sequence, a PCR based approach was used. This involved the use of the polymerase chain reaction to amplify



the precise sequence desired for the expression analysis. To perform the necessary PCR amplification, two oligonucleotide primers were synthesized (Beckman Oligo 1000M DNA synthesizer) have the following sequences:

5 5' primer: 5'TATCTAGACTCAAGCATACGGACAAGGGT 3' (SJ-634) (SEQ.ID.NO.:18)

The italicized bases correspond to nucleotide positions 1 to 21 in the sequence reported in Figure 4. The additional nucleotides 5' of this sequence in the primer are not identical to the promoter sequence, but were included in order to place a XbaI site at the 5' end of the amplification product. The XbaI (5'-TCTAGA-3') (SEQ.ID.NO.:19) site is underlined.

A second (3') primer was synthesized which had the following sequence:

3' primer 5'GGTTATCATTGTATGAACTGA3' (SJ-618) (SEQ.ID.NO.:20)

This primer contains the precise complement (shown in italics) to the sequence reported in Figure 4 from bases 2343 to 2363. This primer was not designed with an additional restriction enzyme site due to the fact that a natural NcoI site (5'-CCATGG-3') (SEQ.ID.NO.:21) straddles the start codon between base pairs 2034 and 2039, thereby allowing for insertion of the storage protein promoter into the appropriate cloning vector.

These two primers were used in a PCR amplification reaction to produce a DNA fragment containing the sequence between nucleotides 1 and 2342 of the flax seed storage protein gene with a XbaI site at the 5' end and a NcoI site 302 base pairs from the 3' end. PCR amplification was performed using the enzyme Pfu (Stratagene) using conditions recommended by the enzyme manufacturer and a temperature program of 94°C (denaturation) for 1 minute, 55°C (annealing) for 1 minute, and 72°C (elongation) for 3.5 minutes. The template was the legumin seed storage protein genomic clone shown in Figure 4.

The resulting amplification product was subsequently digested with XbaI and NcoI to remove the desired 2 kb promoter region. This promoter fragment was cloned into the XbaI and NcoI sites of a XbaI and NcoI digested plasmid designated pGUS1318 (Plasmid pGUSN358S

(Clontech Laboratories) was cut with NcoI and EcoRI and the GUS insert was cloned into pBluescriptKS+ (Stratagene) which was adapted to contain an NcoI site in the multiple cloning site.) The resulting plasmid containing the promoter-GUS fusion was called pPGUS1318. The terminator of the legumin seed storage protein from flax was also amplified from the above mentioned genomic clone. To perform the necessary PCR amplification, oligonucleotide primers were synthesized having the following sequences:

5' primer: 5' GCAAGCTTAATGTGACGGTGAAATAATAACGG 3' (SJ620) (SEQ.ID.NO.:22)

The italicized bases correspond to nucleotide positions 3780 to 3803 in the sequence reported in Figure 4. The additional nucleotides 5' of this sequence in the primer are not identical to the promoter sequence, but were included in order to place a HindIII site at the 5' end of the amplification product. The HindIII site (5'-AAGCTT-3') (SEQ.ID.NO.:23) is underlined.

A second (3') primer was synthesized which had the following sequence:

3' primer 5'TAGGTACCTGGCAGGTAAAGACTCTGCTC3' (SJ-618) (SEQ.ID.NO.:24)

This primer contains the precise complement (shown in italics) to the sequence reported in Figure 4 from bases 4311 to 4290. The additional nucleotides 5' of this sequence in the primer are not identical to the promoter sequence, but were included in order to place a KpnI site at the 5' end of the amplification product. The KpnI site (5'-GGTACC-3') (SEQ.ID.NO.:25) is underlined.

These two primers were used in a PCR amplification reaction to produce a DNA fragment containing the sequence between nucleotides 3779 and 4311 of the flax seed storage protein gene terminator with a HindIII site at the 5' end and a KpnI site at 3' end. Amplification using PCT was as described above. The above pPGUS1318 vector that contains the amplified promoter was digested with XhoI and treated with Klenow to create a blunt end. The vector was subsequently digested with KpnI and the above amplified terminator sequence was inserted so that it was

located 3' of the GUS coding sequence. The resulting vector containing the flax seed storage protein promoter, GUS and the flax seed storage protein terminator is referred to as pPGUST.

The XbaI-KpnI insert of pPGUST which contains the linin promoter-GUS coding sequence-linin terminator sequence was ligated into the XbaI-KpnI sites of pSBS3000 (This vector is a derivative from the *Agrobacterium* binary plasmid pPZP221 (Hajdukiewicz et al., 1994, Plant Molec. Biol. 25: 989-994). In pSBS3000 the plant gentamycin resistance gene of pPZP221 was replaced with parsley ubiquitin promoter-phosphinothricin acetyl transferase gene-parsley ubiquitin termination sequence to confer resistance to the herbicide glufosinate ammonium). The resulting vector is called pSBS2089. In addition the XbaI-KpnI insert of pPGUST which contains the linin promoter-GUS coding sequence-linin terminator sequence was ligated into the XbaI-KpnI sites of the *Agrobacterium* binary plasmid pCGN1559 (MacBride and Summerfield, 1990, Plant Molec. Biol. 14 269-276, confers resistance to the antibiotic kanamycin)). The resulting vector was called pSBS2083. Plasmids pSBS2089 and pSBS2083 were electroporated into *Agrobacterium* strain EHA101. *Agrobacterium* strain EHA101 (pSBS2089) was used to transform flax and *Arabidopsis*, *Agrobacterium* strain EHA101 (pSBS2083) was used to transform *Brassica napus*. Flax transformation was performed essentially as described in Jordan and McHughen (1988) Plant cell reports 7: 281-284, except transgenic shoots were selected on 10  $\mu$ M L-phosphinothricine instead of kanamycin. *Arabidopsis* transformation was done essentially as described in "Arabidopsis Protocols; Methods in Molecular Biology" Vol 82. Edited by Martinez-Zapater JM and Salinas J. ISBN 0-89603-391-0 pg 259-266 (1998) except the putative transgenic plants were selected on agarose plates containing 80 $\mu$ M L-phosphinothricine. *Brassica napus* transformation was done essentially as described in Moloney et al. (1989). Plant Cell Reports. 8: 238-242.

Figure 10 shows the tissue-specific expression of GUS in transgenic flax plants transformed with a linin-GUS gene construct

(pSBS2089). GUS expression was measured in roots (R), stems (S), leaves (L), Buds (B) and embryo (E). Some expression was seen in buds, and maximal expression was achieved in embryo tissues. No detectable expression was seen in any of the untransformed (WT) tissues.

5               Figure 11 shows the temporal expression of GUS in transgenic flax plants transformed with a linin-GUS gene construct (pSBS2089). As can be seen, maximum expression is achieved in mature (pre-dessicated) flax embryos.

10              Figure 12 shows the absolute expression of GUS in transgenic *Brassica napus* plants (L1 to L9) transformed with a linin-GUS gene construct (pSBS2083). As can be seen high level expression can be achieved in *Brassica napus* plants. When comparing individual transgenic plants, a typical variation in expression due to position effect can also be seen.

15              Figure 13 shows expression of GUS in transgenic *Arabidopsis* siliques (transformed with a linin-GUS gene construct (pSBS2089)) during seed development. As can be seen high level expression can also be achieved in *Arabidopsis* seed tissues. Maximum expression is achieved at stage 4 (mature but not fully dessicated) of seed development. No  
20   detectable expression is observed in non-seed tissues such as leaves, stems, roots and silique walls (results not shown).

              While the present invention has been described with reference to what are presently considered to be the preferred examples, it is to be understood that the invention is not limited to the disclosed examples. To  
25   the contrary, the invention is intended to cover various modifications and equivalent arrangements included within the spirit and scope of the appended claims.

              All publications, patents and patent applications are herein incorporated by reference in their entirety to the same extent as if each  
30   individual publication, patent or patent application was specifically and individually indicated to be incorporated by reference in its entirety.

**We Claim:**

1. A method for the expression of a nucleic acid sequence of interest in flax seeds comprising:

5 (a) preparing a chimeric nucleic acid construct comprising in the 5' to 3' direction of transcription as operably linked components:

(1) a seed-specific promoter obtained from flax; and  
10 (2) said nucleic acid sequence of interest wherein said nucleic acid of interest is non-native to said seed-specific promoter;

(b) introducing said chimeric nucleic acid construct into a flax plant cell; and

(c) growing said flax plant cell into a mature flax plant capable of setting seed

15 wherein said nucleic acid sequence of interest is expressed in the seed under the control of said seed-specific promoter.

2. The method according to claim 1 wherein at least one expression characteristic conferred by said seed-specific promoter to its native nucleic acid sequence is conferred to said non-native nucleic acid  
20 sequence.

3. The method according to claim 2 wherein said expression characteristic is timing of expression, level of expression, response to a change in lighting conditions, response to a change in temperature, response to a change in concentration of a chemical agent.

25 4. The method according to claim 1 wherein said flax seed-specific promoter is selected from the group of promoters comprising, oleosin promoters, 2S storage protein promoters and legumin-like seed-storage protein promoters.

005230" E6554960

5. The method according to claim 1 wherein said flax seed-specific promoter comprises:

- 5 (a) a nucleic acid sequence as shown in Figure 1 (SEQ.ID.NO.:1), Figure 2 (SEQ.ID.NO.:4), Figure 3 (SEQ.ID.NO.:6) or Figure 4 (SEQ.ID.NO.:8) wherein T can also be U;
- (b) a nucleic acid sequence that is complimentary to a nucleic acid sequence of (a);
- 10 (c) a nucleic acid sequence that has substantial sequence homology to a nucleic acid sequence of (a) or (b);
- (d) a nucleic acid sequence that is an analog of a nucleic acid sequence of (a), (b) or (c); or
- 15 (e) a nucleic acid sequence that hybridizes to a nucleic acid sequence of (a), (b), (c) or (d) under stringent hybridization conditions.

6. The method according to claim 1 wherein expression of said nucleic acid sequence of interest results in an alteration in protein or fatty acid composition in said seed.

20 7. Transgenic flax seed prepared according to a method comprising:

- (a) preparing a chimeric nucleic acid construct comprising in the 5' to 3' direction of transcription as operably linked components:
- 25 (1) a seed-specific promoter obtained from flax; and
- (2) a nucleic acid sequence of interest wherein said nucleic acid of interest is non-native to said seed-specific promoter;
- (b) introducing said chimeric nucleic acid construct into a flax plant cell; and

- (c) growing said flax plant cell into a mature flax plant capable of setting seed

wherein said nucleic acid sequence of interest is expressed in the seed under the control of said seed-specific promoter.

- 5 8. Transgenic flax seed according to claim 7 wherein at least one expression characteristic conferred by said seed-specific promoter to its native nucleic acid sequence is conferred to said non-native nucleic acid sequence.

9. The method according to claim 8 wherein said expression  
10 characteristic is timing of expression or level of expression.

10. Transgenic flax seed according to claim 8 wherein said seed-specific promoter is a seed storage protein promoter, an oleosin promoter, a 2S storage protein promoter or a legumin-like seed-storage protein promoter.

- 15 11. Transgenic flax seed according to claim 8 wherein said seed specific promoter comprises:

- (a) a nucleic acid sequence as shown in Figure 1 (SEQ.ID.NO.:1), Figure 2 (SEQ.ID.NO.:4), Figure 3 (SEQ.ID.NO.:6) or Figure 4 (SEQ.ID.NO.:8) wherein T can  
20 also be U;
- (b) a nucleic acid sequence that is complimentary to a nucleic acid sequence of (a);
- (c) a nucleic acid sequence that has substantial sequence  
25 homology to a nucleic acid sequence of (a) or (b);
- (d) a nucleic acid sequence that is an analog of a nucleic acid sequence of (a), (b) or (c); or

- (e) a nucleic acid sequence that hybridizes to a nucleic acid sequence of (a), (b), (c) or (d) under stringent hybridization conditions.

12. Transgenic flax seed according to claim 8 wherein expression of said non-native gene of interest results in an alteration in the seed protein or fatty acid composition.

13. Transgenic flax plants capable of setting seed prepared by a method a method comprising:

- (a) preparing a chimeric nucleic acid construct comprising in the 5' to 3' direction of transcription as operably linked components:
  - (1) a seed-specific promoter obtained from flax; and
  - (2) a nucleic acid sequence of interest wherein said nucleic acid of interest is non-native to said seed-specific promoter;
- (b) introducing said chimeric nucleic acid construct into a flax plant cell; and
- (c) growing said flax plant cell into a mature flax plant capable of setting seed

wherein said nucleic acid sequence of interest is expressed in the seed under the control of said seed-specific promoter.

14. An isolated nucleic acid sequence capable of directing seed-specific expression in a plant comprising:

- (a) a nucleic acid sequence as shown in Figure 1 (SEQ.ID.NO.:1), Figure 2 (SEQ.ID.NO.:4), Figure 3 (SEQ.ID.NO.:6) or Figure 4 (SEQ.ID.NO.:8) wherein T can also be U;
- (b) a nucleic acid sequence that is complimentary to a nucleic acid sequence of (a);



- 005280" 6554360
- 5 (c) a nucleic acid sequence that has substantial sequence homology to a nucleic acid sequence of (a) or (b); or
- (d) a nucleic acid sequence that is an analog of a nucleic acid sequence of (a), (b) or (c); or
- 5 (e) a nucleic acid sequence that hybridizes to a nucleic acid sequence of (a), (b), (c) or (d) under stringent hybridization conditions.
15. An isolated chimeric nucleic acid sequence comprising:
- 10 (a) a first nucleic acid sequence comprising a seed-specific promoter obtained from flax which comprises:
- (1) a nucleic acid sequence as shown in Figure 1 (SEQ.ID.NO.:1), Figure 2 (SEQ.ID.NO.:4), Figure 3 (SEQ.ID.NO.:6) or Figure 4 (SEQ.ID.NO.:8) wherein T can also be U;
- 15 (2) a nucleic acid sequence that hybridizes to a nucleic acid sequence of (a) under stringent hybridization conditions;
- (3) a nucleic acid sequence that is complimentary to a nucleic acid sequence of (a); or
- 20 (4) a nucleic acid sequence that has substantial sequence homology to a nucleic acid sequence of (a); and
- (b) a second nucleic acid sequence non-native to said flax seed-specific promoter.
16. A method for the expression of a nucleic acid sequence of interest in a plant seed comprising:
- 25 (a) introducing the chimeric nucleic acid sequence according to claim 15 into a plant cell; and
- (b) growing said plant cell into a mature plant capable of setting seed,

wherein the second nucleic acid sequence is expressed in the seed under the control of the seed specific promoter.

17. A method according to claim 16 wherein said plant cell is selected from the group of plants consisting of soybean (*Glycine max*), rapeseed (*Brassica napus*, *Brassica campestris*), sunflower (*Helianthus annuus*), cotton (*Gossypium hirsutum*), corn (*Zea mays*), tobacco (*Nicotiana tabacum*), alfalafa (*Medicago sativa*), wheat (*Triticum sp.*), barley (*Hordeum vulgare*), oats (*Avena sativa* L.), sorghum (*Sorghum bicolor*), *Arabidopsis thaliana*, potato (*Solanum sp.*), flax/linseed (*Linum usitatissimum*), safflower (*Carthamus tinctorius*), oil palm (*Eleais guineensis*), groundnut (*Arachis hypogaea*), Brazil nut (*Bertholletia excelsa*) coconut (*Cocus nucifera*), castor (*Ricinus communis*), coriander (*Coriandrum sativum*), squash (*Cucurbita maxima*), jojoba (*Simmondsia chinensis*) and rice (*Oryza sativa*).
18. A plant prepared according to the method of claim 16.
19. A plant cell comprising the chimeric nucleic acid sequence according to claim 15.
20. Plant seed comprising the chimeric nucleic acid sequence according to claim 15.
21. Plant seed obtained from a plant prepared according to the method of claim 16.
22. A recombinant expression vector comprising a nucleic acid sequence according to claim 14.
23. A recombinant expression vector comprising a nucleic acid sequence according to claim 15.

Novel methods for the expression of non-native genes in flax seeds and the seeds of other plant species are provided. The methods involve the use of seed-specific promoters obtained from flax. Additionally provided are novel flax seed-specific promoters, chimeric nucleic acid constructs comprising novel flax seed-specific promoters, transgenic plant cells, transgenic plants and transgenic plant seeds containing novel flax seed-specific promoters. The promoters and methods are useful, for example, for altering the seed oil and protein composition in flax seed or other plant seeds

FIGURE 1-1

1 ttcaaaacccgattcccagaggccctattgaagatatgggggaagttcgacgagatcgatgctgggtcgagtgctatg 80  
81 gtgatggtccggtttgggggaggatgagcgagatagccaagactagcattccgttcccacacagagttgggaattgta 160  
161 ccaaataccaaacttgtcgtattggagcgacgatagggacgcggaaaaaacacatccgttgatcagggagttgtacgatg 240  
241 atctcgagccttatgtgtcgaaatccgagggtatgcttacgtgaactacagggatctcgacatcgggatgaatggagga 320  
321 ggtgaaggggatgagaagggtacttatggtgaggctaagggtggtgggggagaagtactttgggggtcaactttgatcgggtt 400  
401 ggttcgggtgaagacgattgttgatcccaataatgtgttttcgaaacgagcagagcattccctcaattccaaactcgggttat 480  
481 aaggatcaatgatcaatgagaatttttcctttcccaatgtgattacaagtctctattgggtcagcttttctcaactgctcctat 560  
561 tcatttagattaattcatacaactattaatttaccagccttttatccggccggttgccgatttattttcttaagtttt 640  
641 agatgaaatgaaaccgatttagtttttattgagatgagattaatctttaatttgcttgaaatttactcacggttgatgtga 720  
721 tatttggaattaaactaaaaatgataaaatcggataaaaaataaaaaatattttaaaaataaaataacataaagaacaata 800  
801 aaataaaataaatttttaatttttcccttggttttctgtgtatcatacatctcttcttacttctttaaaggctt 880  
881 ttcaattatcacttaattaaatacaatagataaaaatcgttaattctataaacatttaacctatacacattgcacggtgaacaat 960  
961 caatatgataataataataataaattcaattatttaattatactacaatttttttaattataaagtttatgcggtcagtt 1040  
1041 tctgcaagctccgagctccttgatcggttagtttctgcggtctcaagggtataacgactcgggagcgagccctttgct 1120  
1121 tccaatggacgggttgcatcttctgcggtcgttgagctcgattggcgtgcatgctggagtcagagttccctacaaaaaac 1200  
1201 cctaaactagaggggtgattaggggtgaaattaggggtgtggcctgggtccattgtccaaaagtttttagtcaacttaaaaaac 1280  
1281 agacttaaaattttatgcttcaaaaatagtttatctgtttattatatattagcgtgtaattagcttggacaaatggggccggcgg 1360

FIGURE 1-2

1361 gtaaggattcgggaccccgatccccgcccatagtgtaatggctcaactgccaagtccaagtcagcattggaccgaaaattattggac 1440  
 1441 acgaagtactaatgtgaaaaaacttttacatttgtttatttttctacttttaatactatatgtctatttttcaaaaatttgaactttta 1520  
 1521 actatgtttttatatagtttagtataatctttaattttttatgcaaaattcatctaattgtattaaaactattttcgatccgtag 1600  
 1601 ctaattatttcgaaggcaagtcaaaagtgttattgttgactatgtgagctaataattgaaccttttatctctcccaaccactc 1680  
 1681 aagttaattgaaccaaactcgatcggtgttggttttcgagctattttcgagccattgttgttatatgcacgtgagatatcaag 1760  
 1761 attgacccgaacacitttattatgataatgtagaaaaagaaaaacatatattctaagactacatgcatgcaaaagtgc aaacccct 1840  
 1841 gcatgaaaagctgctcaacacggtggcatagactcccgccacgtgtccattccacctcatcaccctcaccctcccccacccgttcac 1920  
 1921 ctcttattatatcacaacaatcaatcaatcctactcctccatactcgaacaaatccgaccaacttatcccaatatttccca <sup>CAT</sup> 2000  
 2001 aacttgattaaatttctcagcaat ATG GAT CAG ACG CAC CAG ACA TAC GCC GGA ACC ACG CAG AAC 2065  
 1 M D Q T H Q T Y A G T T Q N 14  
 2066 CCG AGC TAT GGC GGC GGC ACA ATG TAC CAG CAG CAG CCG AGG TCT TAC CAG GCG 2125  
 15 P S Y G G G T M Y Q Q Q Q P R S Y Q A 34  
 2126 GTG AAG GCG GCC ACT GCA GCC ACC GCG GGT GGA TCC CTC ATC GTT CTG TCC GGT CTC ATC 2185  
 35 V K A A T A A T A G G S L I V L S G L I 54  
 2186 CTT ACG GCC ACC GTC ATT TCA CTC ATC ATA GCC ACC CCT CTC CTT GTC ATC TTC AGC CCT 2245  
 55 L T A T V I S L I I A T P L L V I F S P 74  
 2246 GTT CTT GTC CCG GCT CTC ATC ACC GTC GGC CTC TTG ATC ACC GGG TTT CTT GCT TCC GGT 2305  
 75 V L V P A L I T V G L L I T G F L A S G 94  
 2306 GGG TTC GGA GTC GCC GTC ACC GTC TTG TCC TGG ATC TAT AG gtatgtataagcttttgactt 2370  
 95 G F G V A A V T V L S W I Y R 109  
 2371 tagtattgttataaaaataacataaagctgatttatgaacatggatctctcccaacaagagttattttaaagtcatttctcggtctg 2450

FIGURE 1-3

2451 actcgatcgggttggttttgagctactcgggtcacaaatggtcgggtcggtcttgatctgttatactaataatttgaagcc 2530  
 2531 tgaagtttcaattgttctgccccaaacttcccactaccttttgagggtgttaagaagccatacaaaaataattatgaatccct 2610  
 2611 cccaacaactcagaactcgagtcagtggtgttgacggttctctataaaacatttcgaaaaatctttgttcaatgaacgtag 2690  
 2691 aaatgaccatgcttgatgattgtgggtctttataag G TAC GTG ACC GGC GGG CAC CCG GCG GGA GGG 2756  
 110 Y V T G G H P A G G 119  
 2757 GAT TCG CTG GAC CAG GCT AGG TCG AAG TCG GCC AAG GCC AGG GAG GTG AAG GAC AGG 2816  
 120 D S L D Q A R S K L A G K A R E V K D R 139  
 2817 GCG TCG GAG TTC GCA CAG CAG CAT GTC ACA GGT GGT CAA CAG ACC TCT TAA agagagtcctct 2879  
 140 A S E F A Q Q H V T G G Q T S \* 156  
 2880 agttaaattggtcttctgtttctgtttctgtggcggttgtaaaactctcttttaagtggtgctgttttctcttctgtctgtg 2959  
 2960 gttgtaagtgaagtgaatcgaagttccaagttggagatgtttgtaacgatgatgttttctaaataatcagagatatataa 3039  
 Poly A signal  
 3040 aagggttgctaatttagtattgcgtctgatctcggaccacaaactcgcaagtaaaattgcagaggatgagttgtacagaaca 3119  
 3120 agcgtgcattgttctggaagttcatctccttggagccgaccttggttgcttgcccaagtccactagacacaatggt 3199  
 3200 acgagttaaagcctctgtcaaacagatcgctctagcgtcccagaaaaacaccagatttttcgaaaaaccatcggggatcaatt 3279  
 3280 ttcgattcaattccgatcttggaaagtacttgaaacagaagcatgatgctaaaaagataatagaaaaatcgaagcctagaaaaag 3359  
 3360 ttgtacagaaaagcaacaagtcaaaaaatatagatcaacttcaaagggttcaaatattacatcttacagacccccaaaaaatgaca 3439  
 3440 gttaacagaagtcgactaaacagaaaaccagcagcttcacctggaatgaaggagcttttgatcaatccatcctagcttcat 3519  
 3520 tcccccttgaaaattgcagacagagctctcatcctgctaaaagctgggttattcttaaccctgcaatcaataagcatga 3599  
 3600 actaacattggacaccttcatcggcggttgctcgaaaaatcagtcgaggggatttacctgtgtgtgtagtaaacctctc 3679

•



**FIGURE 2-1**

1 tctagacatttgacataaaacccaattc<sup>R1</sup>aaagaacacacaacattgactaacacccaaaaaagaaatagatgtagtgaatttggg 80  
81 agattaaaaaatagaaaacaaactgattcttagaaaagaagagatgattagtgctttcagttcggctgtcaggaatcga 160  
161 gatgttcacttattacattgtcgattc<sup>R2</sup>atctcccaattgtcctggttcctttactgtccgacgcttttttgaatcccag 240  
241 ttaattcccacatcaagtcttccttcagctgcgtagcactgctagctccaacatggagcgtggagtctactcgttcattgggg 320  
321 catcgcaagggttgccttcattgttctgtaccagccagcccccacccctcttggttgtgtggacaattgcggtgaagc 400  
401 gcgcaagttgacatcccatagtctcgacacttcaccatatggatgttttaaaacgtatatcacgagtgcgatctacatgtc 480  
481 ccatcacacacatataaagcaatagtttgggagcttttccatatattgaaacgggcattgacgacttgcctctcgataat 560  
561 ttaatcttttttctcttcagctgattgtgtgcattccattcgggctcagaagcacatcaaaagggatctctccatcgtagt 640  
641 attgggtcgtgtgatgatacgaagcagtcgatgaagtctcctaattgtgcgagctacaggtccgcaaaagacccgcga 720  
721 ggtagatcgtatgctagtagtaccacaaaaatcagtttgtgtagcgggaatcaacactagagactcacccataatgcattctcatg 800  
801 tgtgatgaacagtttatcatttgtgagtcttaggggtcattgtc<sup>R3</sup>gatgacccaatgcacattgagcttatgatagaaatttg 880  
881 aataggaagcgttttccaccagatcacgaatagctaccccttttttcgggcgcaaatctccggcatcctatcttccacc 960  
961 acaacttaagatcgatcggttaaggaaactcaccgaccacacacatcgaataatcttcggtgaccggttccctgttgatca 1040  
1041 agtccctcaatttccctcaacctagtcttcaatcgccgtagcgttatcccccgcatatggactttcatagcgggagcgt 1120  
1121 agccggagacgacgagcaagaaggatgagcggcggcagattgcggctaagaacagagcttccctgccttgctctatggag 1200  
1201 gcagatttctgagttgattggtgatggatttgtgatgtggacacttttaatttaagttgatttttttagcacttcattcacg 1280  
1281 taattaaaaataaatttccagtatatttatatttttcccttacgttatctaattttttgaaaagattaaaaactttgatat 1360  
<sup>R4</sup>



**FIGURE 2-2**

1361 aggaagatcatgacacgctggaagttaagtgaatgagactcctaacaaggtaataacaaaagcagtttcataaaacccgaatga 1440  
R1  
1441 ccttgatcttttactaagcttgagatcattgaacataataattaaatcgtttaatgaaagataaagaacttttaataataaaat 1520  
R4  
1521 cattcaaaaacgagaactgataacaaaaaagcaaaacgcccacaaaataataagacggtggaaggatgatgcagagcc 1600  
R5  
1601 atccaccctttttccaggtttccttactgcttacttctctatgcataatcacaaagacgcccttgaaaacttgttagtcatg 1680  
1681 cagagcccttactgcccaggtcaccgcaccacgtgttactctctatcacttctcctcccttctcctttaagaaccaccacgc 1760  
R5  
1761 cacctccctctcaaaaacactcataaaaaaacccctcttgcatttctcccaagttcaaatagttcacagctaagcaag 1840  
1841 aactcaacaaca ATG GCG GAT CGT ACA ACA CAG CCA CAC CAA GTC CAG GTC CAC ACC CAG CAC 1903  
1 1841 M A D R T T Q P H Q V Q V H T Q H 17  
1904 CAC TAT CCC ACC GGC GGC GCT TTC GGC CGT TAT GAA GGT GGA CTC AAA GGC GGT CCA CAT 1963  
18 H Y P T G G A F G R Y E G G L K G G P H 37  
1964 CAC CAG CAA GGA TCA GGC AGC GGC CCA TCA GCT TCC AAG GTG TTA GCA GTC ATG ACC GCG 2023  
38 H Q G S G S G P S A S K V L A V M T A 57  
2024 CTC CCC ATC GGC GGC ACC CTC CTT GCC TTG GCC GGG ATA ACC TTG GCT GGG ACG ATG ATC 2083  
58 L P I G G T L L A L A G I T L A G T M I 77  
2084 GGG CTG GCG ATC ACC ACC CCG ATT TTT GTC ATC TGC AGC CCT GTT CTA GTC CCG GCC GCT 2143  
78 G L A I T T P I F V I C S P V L V P A A 97  
2144 CTG CTC ATC GGC TTT GCC GTG AGC GCG TTT CTG GCC TCG GGG ATG GCC GGG CTG ACA GGG 2203  
98 L I I G F A V S A F L A S G M A G L T G 117  
2204 CTG ACC TCG CTG TCG TGG TTT GCG AGG TAT CTG CAG CAG GCT GGG CAG GGA GTT GGA GTG 2263  
118 L T S L S W F A R Y L Q Q A G Q G V G V 137  
2264 GGG GTG CCG GAT AGT TTC GAG CAG GCG AAG AGG CGC ATG CAG GAT GCT GCT GGG TAT ATG 2323  
138 G V P D S F E Q A K R R M Q D A A G Y M 157



FIGURE 3-1

```

1  tccactatgtaggtcatatccatcatttttaatttttgggcaccattcaattcccatcttgccttttagggatgtgaatatga 80
   5' primer (1)      AT rich
81  acggccaaggtaaagaaaaataatccaaattaaagcaagagagccaaagtaagataatccaaatgtacacttgtca 160
                        AT rich
161 tcgccgaaattagtaaaatacgcggcatatattgtattccacacattattaaaaataccgtatatgtattggctgcatttgc 240
241 atgaataatactacgtgtaaagcccaaaagaacccacgtgtagcccatgcaaaagttaacactcagacccccattcctcagt 320
   RY                                G box  seed-specific
321 ctccactatataaaacccaccatcccccaatcttaccaaaacccaccacagactcacaaactcgactctcacaccttaaaaga 400
                        TATA
401 ccaatcaccacccaaaaaATGGCAAAGCTGATGAGCCTAGCAGCCGTAGCAACGCAGTTCCTCTTCCTGATCGTGGTGGAC 480
   1      M A K L M S L A A V A T Q F L F L I V V D 21
481 GCATCCGTCGGAACCAACACAGTGATTATCGACGAGGAGACCAACAAAGCCGGGTGGAGGCAAGGTGGCAGGGACAGCAGC 560
22  A S V R T T V I I D E E T N Q G R G G K V A G T A A 48
561 AGTCTGCAGCAGCAGATCCAGCAGCGAGACTTCCTGAGGAGTGCCAGCAGTTTCATGTGGGAGAAAAGTCCAGAGGGGCG 640
49  V C E Q Q I Q Q R D F L R S C Q Q F M W E K V Q R G G 75
641 GCCACAGCCACTATTACAACCAGGGCCGTGGAGGCGGAACAGAGCCAGTACTTCGAACAGCTGTTTGTGACGACCTTA 720
76  H S H Y Y N Q G R G G G E Q S Q Y F E Q L F V T T L 101
721 AGCAATTGCGACCGCGGTGCACCATGCCAGGGGACTTGAAGCGTGCCATCGGCCAAATGAGGCAGGAAATCCAGCAGCA 800
102 S N C A P R C T M P G D L K R A I G Q M R Q E I Q Q Q 128
801 GGGACAGCAGCGGGACAGCAGGAAGTTTCAGAGGTGGATCCAGCAAGCTAAACAAATCGCTAAGGACCTCCCCGGAC 880
129 G Q Q Q Q G Q Q Q Q E V Q R W I Q Q A K Q I A K D L P G Q 155

```

1

FIGURE 4-1

10 20 30 40 50 60 70 80 90 100  
 ctcaagcatacggacaagggtaaataacatagtcaccagaacataataacaaaaagtcagaaagcaagataaaaaaattagctatggacattcaggttc  
 110 120 130 140 150 160 170 180 190 200  
 atattgaaaacatcattatccttagtctgtgaccatccttctcctgctctagttgagaggccttgggactaacgagaggtcagttgggatagcagatcc  
 210 220 230 240 250 260 270 280 290 300  
 ttatcctggactagcctttctgtgttcagagtcttcgtgccgctctacatctatctccattaggtctgaagatgactcttcacaccaacgacgttt  
 310 320 330 340 350 360 370 380 390 400  
 aaggtctctactccttagcttgcaataacctgggcttgcaatacctggagcatcgtgcacgatgattggatactgtggaggaggagtgtttgctgatt  
 410 420 430 440 450 460 470 480 490 500  
 tagagctcccggttggtgatttgacttcgatttcagtttaggctgttgaaaatttttcaggttccattgtgaagccttttagagcttgagcttccctoca  
 510 520 530 540 550 560 570 580 590 600  
 tgttaatgccttgatcgaattctcctagagaaaaagggaagtcgatctctgagtattgaaaatcgaaagtgacacatttttttcaacgtgtccaatcaatcca  
 610 620 630 640 650 660 670 680 690 700  
 caaacaagcagaagacaggtaattcttatactgacaagtaatagtcttaccgtcatgcataataacgtctcgttccgttcaagaggggttttc  
 710 720 730 740 750 760 770 780 790 800  
 cgacatccataacgacccgaagcctcatgaaagcattagggaagaacttttgggttcttcttgcattggcctttataggtgtcagcggagctcgccaattc  
 810 820 830 840 850 860 870 880 890 900  
 ccgtccgactggctccgcaaaaatttcgaacggcaagttatggacttgcaaccataactccacggtattgagcaggacctattgtgaagactcatctcat  
 910 920 930 940 950 960 970 980 990 1000  
 ggagcttcagaatgtggtgtcagcaaaaccaatgaccgaaatccatcacatgacggacgtccagtggtgagcgaaacgaaacaggaagcgccatcttt  
 1010 1020 1030 1040 1050 1060 1070 1080 1090 1100  
 cagagtcgtgagctccacacggattccggcaactacgtgttggcaggcttcgcgtattagagatatgttgaggcaagaccatctgtgccactcgta  
 1110 1120 1130 1140 1150 1160 1170 1180 1190 1200  
 caattacgagagttgtgttttttctcctaagttttctcgttgatggtgagctcatattctacatcgatatggtctctcaacgtcgtttcctctgtcat

**FIGURE 4-2**

1210	1220	1230	1240	1250	1260	1270	1280	1290	1300
ctgatatcccgctcaattgcatccaactgcgcgccgtcccgtagtgccagtccctagtggtcatgcacgcccacaaattggtggtggtgcgggctgcoctgtgctt									
ABRE									
1310	1320	1330	1340	1350	1360	1370	1380	1390	1400
cttaccgatgggtggaggttgagtttgggggtctccgcggcgatgtagtggtgggtgacgggttgggtggtgggtgacggcattgatcaatttacttcttgc									
R1									
1410	1420	1430	1440	1450	1460	1470	1480	1490	1500
ttcaaatctttggcagaaaaaattcattagattagaactggaaccagagtgatgagacggattaaagtcagattccacagagttacatctctttaaga									
R2									
1510	1520	1530	1540	1550	1560	1570	1580	1590	1600
aataatgtaaccccttagactttatataatttgcaattaaaaataatttaaccttttagactttatatataagttttaataactaaagtttaaccactcta									
R2									
1610	1620	1630	1640	1650	1660	1670	1680	1690	1700
ttatttatatcgaaaactattttgtatgtctccctctaaataaaacttggtattgtgtttacagaaacctataatcaaaataatcaataactcaactgaagtttg									
1710	1720	1730	1740	1750	1760	1770	1780	1790	1800
tgcagttaattgaagggttaaacggccaaaatgcactagttattatcaaccgaatagattcacactagatggccattttccatcaatatcatcgccgttctt									
1810	1820	1830	1840	1850	1860	1870	1880	1890	1900
ctttctgtccacatatccctctgaaacttgagagacacctgcacttcattgtcctttattacgtgttacaaaaatgaaacccatgcatccatgcacaaactgaa									
Legumin Vicilin									
1910	1920	1930	1940	1950	1960	1970	1980	1990	2000
gaatggcgcaagaaccttccctccatttcttatgtggcgaccatccatttcacccatctcccgctataaaacaccccccatcacttcacctagaacatca									
CAAT									
2010	2020	2030	2040	2050	2060	2070	2080	2090	2100
tcactacttgcttatccatccaaaagataccccaccatGGCTAGATCATCAAGCCCTTTGCTTCTCTCACTCTGCATTTTCGCCATTCTCTCCACTCTTTC									
TATA									
Signal sequence									
2110	2120	2130	2140	2150	2160	2170	2180	2190	2200
TCTGGGTAGGCAGCAATTCCAGCAGGGGAACGAGTGCAGATCGACAGGATCGACGATCCGAGCCGGACAAACCATCCACGCAGAAAGCTGGCACCATTC									
L G R Q Q F Q Q G N E C Q I D R I D A S E P D K T I Q A E A G T I									
2210	2220	2230	2240	2250	2260	2270	2280	2290	2300
GAGGTATGGGACCGACGACCGCAGCAATTCAGTGCCTGGTGTTCGGCTGTAAAGCGCACCATTTGAGCCCAAAAGGTCTTCTTTCCTTTCATACAGCA									
E V W D Q N R Q Q F Q Q C A G V A V R R T I E P K G L L L P F Y S									

FIGURE 4-3

2310 2320 2330 2340 2350 2360 2370 2380 2390 2400  
ACACCCCTCAGCTCATCTACATCGTTCAAGgtataaaatcagttcatacaatgataaccaccacttcgaatgtatttatcaaatatcaatgatcga  
N T P Q L I Y I V Q

2410 2420 2430 2440 2450 2460 2470 2480 2490 2500  
tgcacctgtatgtgtgtatattcagTAGGGGAGTTACAGGAATCATGTTCCCKAGATGTCCAGAGACATTCGAGGAATCCCAGCAGCAGGACAAC  
G R G V T G I M F P X C P E T F E E S Q Q Q G Q

2510 2520 2530 2540 2550 2560 2570 2580 2590 2600  
AGGGCCAAACAGGGTAGTTCCCAAGACACGACACAGATCCGCGCTTCGGTGAAGGTGACGTCAATTGCCGTCCCTGCCGGTGTAGCCCACTGGTCCTA  
Q G Q Q G S S Q D Q H Q K I R F R E G D V I A V P A G V A H W S Y

2610 2620 2630 2640 2650 2660 2670 2680 2690 2700  
CAACGATGGCAACGAACAGTCTATGGCCATTGTTGTCCATGACACTTCCAGCCACCTCAACCAACTGGACAACAACCCAGGgtatataagcattgcccgt  
N D G N E P V M A I V V H D T S S H L N Q L D N N P R

2710 2720 2730 2740 2750 2760 2770 2780 2790 2800  
agttgctaataaattgcacacaattggaaactctattttcagtatctataaacttttctctttttggcagAACTTCTACTTGGCAGGAAACCCGAGAGAC  
N F Y L A G N P R D

2810 2820 2830 2840 2850 2860 2870 2880 2890 2900  
GAGTTCGAACAATCGCAGCAAGGAGGCGAGCTGAGCCGTGGGGAGAGTGAAGGTGGACGAGGACGGGAACCTCTTCAACCTGCAACAACCTCTTCTT  
E F E Q S Q Q G G R L S R G E S E G G R G R E P L Q P A T T S S

2910 2920 2930 2940 2950 2960 2970 2980 2990 3000  
GCGGAATCGACTCCAAGCTCATCGCGAGGCGTTCAATGTGACGAGAACGTGGCAAGGAGGCTACAGCGGAGAACGACAACAGAGGCCAGATCGTCCG  
C G I D S K L I A E A F N V D E N V A R R L Q S E N D N R G Q I V R

3010 3020 3030 3040 3050 3060 3070 3080 3090 3100  
AGTCGAAGCGGAGCTCGACATCGTCAGACCTCCGACCACTATCCAGGAGGAGTCAAGGAGGAGGAGTGGTGGTGGCGCTACTACTCCAATGGA  
V E G E L D I V R P P T S I Q E E S Q E Q G G R G G R Y Y S N G

3110 3120 3130 3140 3150 3160 3170 3180 3190 3200  
GTGAGGAGACCTTCTGCTCCATGAGACTAATTGAGAACATCGGCGATCCTTCTCGGGCAGACATTTTCTACTCCAGAAGCCGGCGGCTTAGATCCCTCA  
V E E T F C S M R L I E N I G D P S R A D I F T P E A G R V R S L

**FIGURE 4-4**

3210 3220 3230 3240 3250 3260 3270 3280 3290 3300  
ACAGCCACAACTCCCGTCCTGCAATGGATCCAGCTTAGCGCCGAGAGAGGGGTTCTCTACAATgtatagatctactcaacgcacaaacttaaaattga  
N S H N L P V L Q W I Q L S A E R G V L Y N

3310 3320 3330 3340 3350 3360 3370 3380 3390 3400  
atccctaattatttaattcaccgatatactgaccgcacgggtttgaatttttagGAAGCGATCAGGCTGCCGCACTGGAACATCAACGCACACAGCATAGT  
E A I R L P H W N I N A H S I V

3410 3420 3430 3440 3450 3460 3470 3480 3490 3500  
GTACGGGATCAGAGGACAAGCCAGAGTCCAGATCGTGAACGAGGAAGGSAATTTCGGTGTTCGATGGAGTCTGCAGGAAGACAGGTGGTGACGGTGCCG  
Y A I R G Q A R V Q I V N E E G N S V F D G V L Q E G Q V V T V P

3510 3520 3530 3540 3550 3560 3570 3580 3590 3600  
CAGAACTTCGGGGTGGTAAAGAGATCCCAGAGCGAGAGGTTTGAGTGGGTGGCGTTCAAGACCAACGACACGCGATGGTGAACCTCGTAGCCGGGAGGA  
Q N F A V V K R S Q S E R F E W V A F K T N D N A M V N S L A G R

3610 3620 3630 3640 3650 3660 3670 3680 3690 3700  
CATCGGACTAAGGGCGATCCCGCGGATGTACTGGCTAACGCCCTGAGGGGTGTCGCCGAGGAGGCGGAGGGTGAAGTTCAACAGGAGGAGACTCA  
T S A V R A I P A D V L A N A W R V S P E E A R R V K F N R Q E T H

3710 3720 3730 3740 3750 3760 3770 3780 3790 3800  
CTTGGCTAGCACCCAGGGGCCAGTCCAGGTCGCCCGGGAGGTTGAATGTCTCAAGGAGGTGATCAACTTGCTTTATGTAaaatgtgacggtgaaataataa  
L A S T R G Q S R S P G R L N V V K E V I N L L M \*

3810 3820 3830 3840 3850 3860 3870 3880 3890 3900  
cggtaaaatatatgtaataataataataaaagccacaaagtgagaatgaggggaagggaatgtgtaatgagccagtagccggtggtgtaattttg

3910 3920 3930 3940 3950 3960 3970 3980 3990 4000  
tatcgtattgtcaataaatcatgaattttgtgtgttttttaaatcatgaatttttaaaattttataaaataatctccaatcgggaagaacaac

4010 4020 4030 4040 4050 4060 4070 4080 4090 4100  
attccatatccatggatgttttttaccctaaatctagttctttgagagatgaagcatcaccgaacagttcttgcaactatccctcaaaagctttaaaatga

4110 4120 4130 4140 4150 4160 4170 4180 4190 4200  
acaacaaggaaacagagaaacgttccaaagatcccaaacgaaacatatattctataactaataattattattactactgccccgggaatcacaaatccct



FIGURE 4-5

4210 4220 4230 4240 4250 4260 4270 4280 4290 4300  
 gaatgattcctattactacaagccttgttggcggcgagaaagtgatcgcgcgagaaagcagcgactcggagacgagccttggatgagcagagtc  
 4310 4320 4330 4340 4350 4360 4370 4380 4390 4400  
 ttacacctgccagggcgtgaaggggaagagcgcccttctggagtaggagttcagcaagcgcggttccttggcggagtaagcggacgtaagggtggnctgtc  
 4410 4420 4430 4440 4450 4460 4470 4480 4490 4500  
 gacgtcntcgttctcnggagggcgnattcatgaagggttaagtcnatctctcgtcgtcagtgagcnaaaagacgttgggaaaccgctcgcncgt  
 4510 4520 4530 4540 4550 4560 4570 4580 4590 4600  
 ttggggcagtcagtcngcggggcagcgttccctcctcgtcgtcctccanaancnangtanatttaaaganaatgggaaattaatgnaatnannaggagg  
 4610 4620 4630 4640 4650 4660 4670 4680 4690 4700  
 attgnaacggtcngancnngnangaaanagtttttannggtttaataactgggggagtnagnccnctggttccngttagangaaaccaagnnccgg  
 4710 4720 4730 4740 4750 4760 4770 4780 4790 4800  
 gaggnnttcannnnnaggagaaaaaggannncatttnannangcngagggacatgaancggtacngagctgnggttcannnnancgcnngnngnagtcc  
 4810 4820 4830 4840 4850 4860 4870 4880 4890 4900  
 cnngggaacnggntggggttnaagggaanggaacattngtngnangganaaanaccnttttaacnattgcctttgcaggnnngtntngcncntncgggt  
 4910 4920 4930 4940 4950 4960 4970 4980 4990  
 nacatncgctgcatgggctttgggngccnanagnagccncanggggnanncngccncttctncaangcgcctnaagttcnattgtanatggncgttg

FIGURE 5




FIGURE 6






FIGURE 7

8 12 16 20 22



10J  
(L-isoform)



3T  
(H-isoform)

**FIGURE 8**






Figure 9.1



005280" 26554960

Figure 9.2



**FIGURE 10**







FIGURE 11



**FIGURE 12**






FIGURE 13



SEQUENCE LISTING

<110> Chaudhary, Sarita  
van Rooijen, Gijs  
Moloney, Maurice  
Singh, Surinder

<120> Flax Seed Specific Promoters

<130> 9369-151

<140>

<141>

<150> 60/151044

<151> 1999-08-27

<150> 60/161,722

<151> 1999-10-27

<160> 25

<170> PatentIn Ver. 2.0

<210> 1

<211> 4305

<212> DNA

<213> Linum usitatissimum

<400> 1

ttcaaaaccc	gattccccgag	gcggccctat	tgaagatatg	ggggaagttc	gacgagatcg	60
atgtcggggtc	gagtgcctatg	gtgatgggtgc	cgtttggggg	gaggatgagc	gagatagcca	120
agactagcat	tccgttccca	cacagagttg	ggaatttgta	ccaaatccaa	cacttgctcg	180
attggagcga	cgatagggac	gcggaaaaac	acatccggtg	gatcagggag	ttgtacgatg	240
atctcgagcc	ttatgtgtcg	aagaatccga	ggtatgctta	cgtgaactac	agggatctcg	300
acatcgggat	gaatggagga	ggtgaagggg	atgagaaggg	tacttatggg	gaggctaagg	360
tgtgggggga	gaagtacttt	ggggctcaact	ttgatcgggt	ggttcgggtg	aagacgattg	420
ttgatcccaa	taatgtgttt	cgaaacgagc	agagcattcc	ctcaattcca	actcggttat	480
aaggatcaat	gatcaatgag	aattttcctt	tccaatgtga	ttacaagttc	tattgggtca	540
gctttctcaa	ctgctcctat	tcatttagat	taattcataa	caactattaa	tttaccagcc	600
ttttatccgg	cccgttggcc	gattttattt	cttaagtttt	agatgaaatg	aaaccgattt	660
agttttttat	gagatgagat	taatcttaat	ttgcttgaaa	tttactcacg	gttgatgtga	720
tatttgggaat	taactaaaat	gataaatatc	ggataaaaaat	aaaaatat	aaaataaata	780
acataaacat	aagaacaata	aaataaataa	atttaatttt	aatttatttc	cttgttttct	840
ttctgtatca	tacatctctt	ctcttacttc	ttaaaggctt	ttcaattatc	acttaattaa	900
atacaataga	taaatcggtta	attctataac	attaacctat	acacttgcac	ggtgaacaat	960
caatatgata	atataataat	aatataataa	ttcaattatt	aatctacaat	tttttaatta	1020
taaagtttat	gcggtcagtt	tctgcaagct	ccgagctcct	tgtcatcggt	agtttctgcg	1080
gtctcaagg	ataacgactc	ggagcgacga	gccctttgct	tccaatggac	gggttgccatt	1140
tctgcccgtcg	ttgagctcga	ttggcgtgtc	atgctggagt	cagagttcct	acaaaaaac	1200
cctaaactag	aggggtgatta	gggtgaaatt	aggggtgttg	cctgggttcc	attgtccaaa	1260
gttttagtca	acttaaaaac	agacttaaat	tttatgcttc	aaaatagttt	atctgttatt	1320
atattagcgt	gtaattagtc	ttgacaatgg	ggcgggacgg	gtacggattc	gggaccccca	1380
tccccgccca	tagtgtaatg	gctcaactgc	caagtcagca	ttggaccgaa	attattggac	1440
acgaagtact	aatgtgaaaa	actttacatt	tgttattttc	tactttaata	ctatgctatt	1500
ttcaaaattt	gaactttaat	actatgtttt	tatatagttt	agtatatctt	aatttttatg	1560
caaattcatc	taattgtatt	aaactatttt	cgatccgtag	ctaattattt	cgaaggcaag	1620
tcaaagtgtt	attgtggact	atgtgagcta	atattgaacc	tttatctctc	ccaaccactc	1680
aagttaattg	aaccaaaactc	gatcggttgg	gtttcgagct	atcttcgagc	attgttggtta	1740
tatgcacgtg	agatatcaag	attgaccoga	acactttatt	atgataatgt	agaaaaagaa	1800
aacatattct	aagactacat	gcatgcaaag	tgcaaccctt	gcatggaaag	ctgctcaaca	1860
cgtggcatag	actcccgcga	cgtgtccatt	ccacctcatc	acctcaccct	caccgttcac	1920
ctcttattat	atcacaacca	tcaattcaatc	ctactctccc	atactcgaac	aaatccgacc	1980
aacttatacc	aattattccca	aacttgatta	atctctcagc	aatatggatc	agacgcacca	2040
gacatacgcc	ggaaccacgc	agaaccgcag	ctatggcggc	gggggcacaa	tgtaccagca	2100

```

gcagcagccg aggtcttacc aggcgggtgaa ggcggccact gcagccaccg cgggtggatc 2160
cctcatcggt ctgtccgggtc tcatccttac ggccaccgtc atttcactca tcatagccac 2220
ccctctcctt gtcactttca gccctgttct tgtoocggct ctcatcacccg tcgggctctt 2280
gatcaccggg tttcttgctt ccgggtgggtt cggagtgcgc gccgtcaccg tcttgctctg 2340
gatctatagg tatgtataag ctttggactt tagtattgtt ataaaataca taagctgatt 2400
tatgaacatg gatctcccaa caagagttat ttaaattgcat tctcgggtctg actcgatcgg 2460
ttgggttttg agctactcgg tcacaatggt cgggtcgggt ctggatctgt tatactaata 2520
tttgggaagc tgaagtttca ttgttctgcc ccaacttccc actacctttt gaggggtgta 2580
agaagccata caaactaatt atgaatccct cccaacaact cagaactcga gtcagtgggt 2640
tgtgacgggt ctctataaac atttcgaaaa tctttgttca atgaacgtag aaatgaccat 2700
gcttgatgat tgtgggtctt ataaggtagc tgaccggcgg gcaccggcg ggaggggatt 2760
cgctggacca ggctagggtc aagctggccg gaaaggccag ggaggtgaag gacaggcgt 2820
cggagtctgc acagcagcat gtcacagggt gtcacacagac ctcttaaaga gagtccctta 2880
gttaaattgg tcttcgtttc tgtttcgtgg cggcttgtaa actctctttt aagtgtgctg 2940
ttttcctttt gtctcgtgtg ttgtaagtga aagtgtaatc gaagttccaa gttggagatg 3000
tttgtaacga tgatgttttc taataatcag agatattaaa agggttgcta atttagtatt 3060
gcgtctgata tcggacacaa ctcgcaagta aaattgcaga ggatgagttg tacagaacaa 3120
gcgtgcattg ttctggaagt tcatctcctt ggagccgacc ttgttgcttg cagtttcgcc 3180
aagtccacta gacaatgtta cgagttaagc ctctgtcaaa cagatcgctc tagcgtccca 3240
gaaaacacca gatttttcga aaacctcgg ggatcaattt tcgattcaat tccgatcttg 3300
gaagtacttg aacagaagca tgatgctaaa agataataga aaatcgaagc ctagaaaagt 3360
tgtacagaaa gcaacaagtc aaaaatatag atcaacttca aaggttcaaa ttacatctta 3420
cagaccccaa aaaatgacag ttaacagaag tcgactaaac agaaaccagc cagcttcacc 3480
tggaatgaag gagctttgat caatccatcc tagcttcatt cccctttgaa attgcagaca 3540
gagctctcat cctgctaaag ctggtggctt attcttaacc ctgcaatcaa taagcatgaa 3600
ctaacattgg acaccttcat cggcggattg ctcgaaaatc agtgagcgag ggatttacct 3660
gtgtgtgtag taacctctct ccttgtacat aaaatctgga aattccggca tcaactactg 3720
ccacctttct gcttaagggtg attttatcac caaggctgag cgtgattcct tgcgtcttgc 3780
tccgaatcct gatgtatcca ctgagctttc catctccttc cttctccagg cttatgttca 3840
ccaatgcgtc ctgcgccgaac acactcttgg cgtacaagtt cgcagccagg aatccacact 3900
ctccatcaag tgcagacctg caaaccacaa ataagaacac aaactccaaa gtcaacgatc 3960
aattctccgc cttttatgaa gaaaaggaaa cttctgggta cttacgggtg cgtcagacac 4020
ttcatatttg tagacttgat gatatggtcc aggaattcct tctcgttctg aattgttggtg 4080
ttaacagcaa cctgacagac agaaagatat cgcaaattta agatactggg atgactaggc 4140
acagagaaat gaaatctaatt tctagaagta aaaccttatt ttccattca aattctgccc 4200
acatagtccg gaacgcagca tccgagcaag aagcaggaga gatgtaatcc atgatatcga 4260
tgtggatatc gttgaggacg acaactgaac gttccatcac attgg 4305

```

<210> 2  
 <211> 109  
 <212> PRT  
 <213> Linum usitatissimum

<400> 2  
 Met Asp Gln Thr His Gln Thr Tyr Ala Gly Thr Thr Gln Asn Pro Ser  
 1 5 10 15  
 Tyr Gly Gly Gly Gly Thr Met Tyr Gln Gln Gln Gln Pro Arg Ser Tyr  
 20 25 30  
 Gln Ala Val Lys Ala Ala Thr Ala Ala Thr Ala Gly Gly Ser Leu Ile  
 35 40 45  
 Val Leu Ser Gly Leu Ile Leu Thr Ala Thr Val Ile Ser Leu Ile Ile  
 50 55 60  
 Ala Thr Pro Leu Leu Val Ile Phe Ser Pro Val Leu Val Pro Ala Leu  
 65 70 75 80  
 Ile Thr Val Gly Leu Leu Ile Thr Gly Phe Leu Ala Ser Gly Gly Phe  
 85 90 95  
 Gly Val Ala Ala Val Thr Val Leu Ser Trp Ile Tyr Arg  
 100 105

00459-02500

<210> 3  
 <211> 46  
 <212> PRT  
 <213> Linum usitatissimum

<400> 3  
 Tyr Val Thr Gly Gly His Pro Ala Gly Gly Asp Ser Leu Asp Gln Ala  
     1                    5                    10                    15  
 Arg Ser Lys Leu Ala Gly Lys Ala Arg Glu Val Lys Asp Arg Ala Ser  
             20                    25                    30  
 Glu Phe Ala Gln Gln His Val Thr Gly Gly Gln Gln Thr Ser  
             35                    40                    45

<210> 4  
 <211> 3501  
 <212> DNA  
 <213> Linum usitatissimum

<400> 4  
 tctagacatt tgacataaac cgaattcaaa gaacacaaca ttgactaaca ccaaaaagaa 60  
 atagagtagt gaaatttgga agattaaaaa atagaaacaa actgattctt agaaagaaga 120  
 gatgattagg tgctttcagt tcggtctgtc aggaaatcga gatgttcaact tatttacatt 180  
 gtgcattcat ctcccaattg tcctggttcc tttactgtcc gacgcttttt tgaatcccag 240  
 ttaattccca tcaagtcttc cttcagctgc gtagcactgc tagctccaac atggagcgtg 300  
 gagtctactc gttcatgggg catcgcaaag gtttgcttc atgttctgct accagccagc 360  
 gccacccgcc tcttggttgt gtggacaatt gcggtgaagc gcgcaagttg acatcccata 420  
 gtctcgacac ttcaccatat ggatgtttaa aacgtatatc acgagtgcga tctacatgtc 480  
 ccatacaccc acatataaag caatagtttg ggagcttttc atatttgaaa cgggcattga 540  
 cgacttgccc tctcgataat ttaatctttt tttctcttca gctgattgtg tgcattccatt 600  
 cgggctcaga agcacatcaa agggatctct ccactgtagt attgggtcgt gtcgtatgat 660  
 acgaagcagt cgatgaagtt tcctaattgt cgagctacag gctccgcaaa gaaccgcga 720  
 ggtagatcgt atgctagtac ccaaaaatca gtttgctgta gcggaatcaa cactagagac 780  
 tcaccctaag gcattctcat tgtgatgaac agtttatcat ttgtgagtct aggggtcatt 840  
 cgcgatgacc caatgcacat tgagcttatg atagaatttg aataggaagc gttttccacc 900  
 cagatcacga atagctaccc ctttttcggg cgccaaattt ccggcatcct atcttccacc 960  
 acaacttaaa gatgcgatcg gtaaggaact caccgaccac acacatcgaa taatcttcgg 1020  
 tgaccgggtc ctggtgatca agtccctcaa tttcctcaac ctagtcttca atcgccgcta 1080  
 gcggtatccc ccgcataatg actttcatag cgcggagcgt agccggagac gacgagcaag 1140  
 aaggatgagc ggcggcagat tgccgctaaa gaaacgagct tcctgccttg ctctatggag 1200  
 gcagatttct gagttgatgg tgatggattt gtgatgtgga cacttttaat ttaagttgat 1260  
 tttttagcac ttcattcacg taattaaata aataatttcc agtattttat atttatttcc 1320  
 ttacgttatc taattttttg aaagattaaa actttgatat aggcaagatc atgacacgtc 1380  
 gaagttaagt gaatgagact cctaacaagg taataacaaa gcagttcata aaccgaatga 1440  
 ccttgatctt tactaagctt gagatcattg aacatataat taaatacgtt aatgaaagat 1500  
 aagaacttta atataaaaaat cattcaaaac gagaaactga taacaaaaac aaagcaaacg 1560  
 gccacaacaaa taatagacgg tggaaggatg atgcagagcc atccaccctt ttttcccagt 1620  
 ttccttactg cttacttctc tatgcataac acaagacgcc cttgaaaact gttagtcatt 1680  
 cagagccctt actcgccagg tcacgcgacc acgtgttact ctatcacttc tcttcccttt 1740  
 cctttaaaga accaccacgc cacttccctc tcacaaacac tcataaaaaa accacctctt 1800  
 gcatttctcc caagttcaaa ttagttcaca gctaagcaag aactcaacaa caatggcgga 1860  
 tcgtacaaca cagccacacc aagtccaggt ccacaccag caccactatc ccaccggcgg 1920  
 ggctttcggc cggttgaag gtggactcaa agcggttcca catcaccagc aaggatcagg 1980  
 cagcggccca tcagcttcca aggtgttagc agtcatgacc gcgtcctcca tcggcggggac 2040  
 cctccttgcc ttggccggga taaccttggtc tgggacgatg atcgggctgg cgatcaccac 2100  
 ccgatttttt gtcattctgca gccctgttct agtcccgcc gctctgctca tcgggttttg 2160  
 cgtgagcgcg tttctggcct cggggatggc cgggctgaca gggctgacct cgctgtcgtg 2220  
 gtttgcgag tatctggcag aggtcgggca gggagtggga gtgggggtgc cggatagttt 2280  
 cgagcaggcg aagaggcgca tgcaggatgc tgctgggtat atggggcaga agaccaagga 2340  
 agttgggcag gagatccaga ggaagtctca ggatgtgaaa gcatcagaca aataaggtga 2400  
 taataagggg ttttgggttc gtgtgtaaag tggtaaaatg gaaattctgg gttttactgg 2460  
 acttttgcag ttagtggaat gaatgagttc ttgttctctt ttgtctttta atcataaagt 2520  
 aagaagcagc atttcatgtt ctggttgaat attgtcaaga attcgcaaca aatttagcta 2580  
 aaccagttca atcttaccgg ttagacgact tcccagtaag aaacattcca ggtccatccc 2640

005280" E554360



```

ctttagggat gtgaatatga acggccaagg taagagaata aaaataatcc aaattaaagc 120
aagagaggcc aagtaagata atccaaatgt acacttggtca tcgccgaaat tagtaaaata 180
cgcgccatat tgtattccca cacattatta aaataaccgta tatgtattgg ctgcatttgc 240
atgaataata ctacgtgtaa gcccaaaaga acccacgtgt agcccatgca aagttaacac 300
tcacgacccc attcctcagt ctccactata taaacccacc atccccaatc ttaccaaac 360
caccacacga ctcaaacctc gactctcaca ccttaaagaa ccaatcacca ccaaaaaatg 420
gcaaagctga tgagcctagc agccgtagca acgcagttcc tcttcctgat cgtgggtggac 480
gcatccgtcc gaaccacagt gattatcgac gaggagacca accaaggccg cgggtggaggc 540
aaggtggcag ggacagcagc agtctgcgag cagcagatcc agcagcgaga cttcctgagg 600
agctgccagc agttcatgtg ggagaaagtc cagaggggag gccacagcca ctattacaac 660
cagggccgtg gaggaggcga acagagccag tacttcgaac agctgtttgt gacgacctta 720
agcaattgag caccgcggtg caccatgccca ggggacttga agcgtgccat cggccaaatg 780
agggcaggaaa tccagcagca gggacagcag cagggacagc agcaggaagt tcagaggtgg 840
atccagcaag ctaaacaat cgctaaggac ctcccggagc agtgccgcac ccagcctagc 900
caatgccagt tccagggcca gcagcaatct gcatggtttt gaaggggtga tcgattatga 960
gatcgtaaaa agacactgct aggtgttaag gatggataat aataataata atgagatgaa 1020
tgtgttttaa gttagtgtaa cagctgtaat aaagagagag agagagagag agagagagag 1080
agagagagag agagagagag agaggctgat gaaatgttat gtatgtttct tggtttttaa 1140
aataaatgaa agcacatgct cgtgtggttc tatcgaatta ttcggcggtt cctgtgggaa 1200
aaagtccaga agggcgggccg cagctactac tacaaccaag gccgtggagg agggcaacag 1260
agccagcact tcgatagctg ctgcatgat cttaagcaat tgaggagcga gtgcacatgc 1320
aggggactgg agcgtgcaat cggccagatg aggcaggaca tccagcagca gggacagcag 1380
caggaagtgg agaggtggtc ccatcaatct aaacaagtcg ctagggacct tccgggacag 1440
tgccgcaccc agcctagccg atgccagctc caggggcagc agcagctctg atggttttga 1500
agtgggtgatc gatgagatcg tataaagaca ctgctagggt ttaaggatgg gataataaga 1560
tgtgttttaa gtcattaacc gtaataaaaa gagagagagg ctgatggaat gttatgtatg 1620
tatgttttct ggttttttaa attaaatgga aagcacatgc tcgtgtgggt tctatc 1676

```

<210> 7  
 <211> 174  
 <212> PRT  
 <213> Linum usitatissimum

<400> 7  
 Met Ala Lys Leu Met Ser Leu Ala Ala Val Ala Thr Gln Phe Leu Phe  
 1 5 10 15  
 Leu Ile Val Val Asp Ala Ser Val Arg Thr Thr Val Ile Ile Asp Glu  
 20 25 30  
 Glu Thr Asn Gln Gly Arg Gly Gly Gly Lys Val Ala Gly Thr Ala Ala  
 35 40 45  
 Val Cys Glu Gln Gln Ile Gln Gln Arg Asp Phe Leu Arg Ser Cys Gln  
 50 55 60  
 Gln Phe Met Trp Glu Lys Val Gln Arg Gly Gly His Ser His Tyr Tyr  
 65 70 75 80  
 Asn Gln Gly Arg Gly Gly Gly Glu Gln Ser Gln Tyr Phe Glu Gln Leu  
 85 90 95  
 Phe Val Thr Thr Leu Ser Asn Cys Ala Pro Arg Cys Thr Met Pro Gly  
 100 105 110  
 Asp Leu Lys Arg Ala Ile Gly Gln Met Arg Gln Glu Ile Gln Gln Gln  
 115 120 125  
 Gly Gln Gln Gln Gly Gln Gln Gln Glu Val Gln Arg Trp Ile Gln Gln  
 130 135 140  
 Ala Lys Gln Ile Ala Lys Asp Leu Pro Gly Gln Cys Arg Thr Gln Pro  
 145 150 155 160  
 Ser Gln Cys Gln Phe Gln Gly Gln Gln Gln Ser Ala Trp Phe  
 165 170

005280 "0554950"



<210> 8  
<211> 4999  
<212> DNA  
<213> Linum usitatissimum

<400> 8  
ctcaagcata cggacaaggg taaataacat agtcaccaga acataataaa caaaaagtgc 60  
agaagcaaga taaaaaaatt agctatggac attcaggttc atattggaaa catcattatc 120  
ctagtcttgt gaccatcctt cctcctgctc tagttgagag gccttggggac taacgagagg 180  
tcagttggga tagcagatcc ttatcctgga ctagcctttc tgggtgtttca gagtcttcgt 240  
gccgccgtct acatctatct ccattaggtc tgaagatgac tcttcacacc aacgacgttt 300  
aagggtctcta tcctactcct agcttgcaat acctggccttg caatacctgg agcatcgtgc 360  
acgatgattg gatactgtgg aggaggagt tttgctgatt tagagctccc ggttgggtga 420  
tttgacttctg atttcagttt aggcctgttg aaatttttca ggttccattg tgaagccttt 480  
agagcttgag cttccttcca tgtaaatgcc ttgatcgaat tctcctagag aaaagggaag 540  
tcgatctctg agtattgaaa tcgaagtga catttttttt caacgtgtcc aatcaatcca 600  
caaacaaagc agaagacagg taatctttca tacttatact gacaagtaat agtcttaccg 660  
tcatgcataa taacgtctcg ttcttccaag aggggttttc cgacatccat aacgacccga 720  
agcctcatga aagcattagg gaagaacttt tgggtctctc tgatcatggcc tttataggtg 780  
tcagccgagc tcgccaatte ccgtccgact ggctccgcaa aatattcgaa cggcaagtta 840  
tggacttgca accataactc cacggtattg agcaggacct attgtgaaga ctcatctcat 900  
ggagcttcag aatgtgggtg tcagcaaacc aatgaccgaa atccatcaca tgacggacgt 960  
ccagtggttg agcgaacga aacaggaagc gcctatcttt cagagtcgtg agctccacac 1020  
cggattccgg caactacgtg ttgggcaggc ttccgcgtat tagagatatg ttgaggcaag 1080  
acccatctgt gccactcgta caattacgag agttgttttt tttgtgattt tctaaagttt 1140  
ctcgttgatg gtgagctcat attctacatc gtatggcttc tcaacgtcgt ttctgtcat 1200  
ctgatatccc gtcatttgca tccacgtgcg ccgctcccg tgccaagtcc ctaggtgtca 1260  
tgcaagccaa attgggtgtg gtgcgggctg ccctgtgctt ctaccgatg ggtggagggt 1320  
gagtttgggg gtctccgcgg cgatggtagt ggggtgacgg tttggtgtgg gttgacggca 1380  
ttgatcaatt tacttcttgc ttcaaattct ttggcagaaa acaattcatt agattagaac 1440  
tggaaaccag agtcatgaga cggattaagt cttatccaa cagagttaca tctcttaaga 1500  
aataatgtaa cccctttaga ctttatatat ttgcaattaa aaaaataatt taacttttag 1560  
actttatata tagttttaat aactaagttt aaccactcta ttatttatat cgaaactatt 1620  
tgtatgtctc cctctaaaat aaacttggtt ttgtgtttac agaactata atcaaataat 1680  
caatactcaa ctgaagtttg tgcagttaat tgaagggatt aacggccaaa atgcactagt 1740  
attatcaacc gaatagattc acactagatg gccatttcca tcaatatcat cgccgttctt 1800  
cttctgtcca catatccct ctgaaacttg agagacacct gcacttcatt gtccattatta 1860  
cgtgttacaa aatgaaacc atgcatccat gcaaactgaa gaatggcgca agaacccttc 1920  
ccctccattt cttatgtggc gaccatccat ttcacctc ctactacttg ttaattccatc caaaagatac ccaccatggc 2040  
tcaattcacc tagaacatca tcactacttg ttctctcact ctgcattttc gccattctct tccactcttc 2100  
tagatcatca agccctttgc agcaggggaa cgagtgccag atcgacagga tcgacgcatc 2160  
tctgggtagg cagcaattcc aggcagaagc tggcaccatc gaggtatggg accagaaccg 2220  
cgagccggac aaaaccatcc gtgttgccgt tgaaggcgc accattgagc ccaaaggctc 2280  
ccagcaattc cagtgcgtg gtgttgccgt gtcactctac atcgttcaag gtataaatta 2340  
tctcttgcc ttctacagca acacccctca cgaatgtatt tatcaaatat caatgatcga 2400  
aatcagttca tacaatgata accaccactt aggggagttt caggaatcat gttcccakga 2460  
tgcacctgta tgtgttgtgt atattcaggt caaggacaac agggccaaca ggtagttcc 2520  
tgtccagaga cattcgagga atcccagcag cgtgaagggt acgtcattgc cgtccctgcc 2580  
caagaccagc accagaagat ccgcccgttc aacgaaccag tcatggccat tgttgtccat 2640  
ggtgtagccc actggtccta caacgatggc aacaacccca gggatatata gcattgccgt 2700  
gacacttcca gccacctcaa caactggac caattggaac tctattttca gtatctaata actttttcct 2760  
agttgctaatt aaattgcaca tggcagggaa cccgagagac gagttcgaac aatcgacga 2820  
tttttggcag aacttctact gggagagtga aggtggagca ggacgcaggg aacctcttca 2880  
aggaggcagg ctgagccgtg gcggaatcga ctccaagctc atcgcgagg agatcgtccg 3000  
acctgcaaca acctcttctt ggctacagag tccgaccagt atccaggagg agtcacagga 3060  
cgacgagaac gtggcaagga tcgtcagacc cccgctacta gtggaggaga cttctgtctc 3120  
agtcaaggc gagctcgaca gccgctacta ttctcgggca gacattttca ctccagaagc 3180  
gcagggaggt cgtgggtgtg attgagaaca cctccccgtc ctgcaatgga tccagcttag 3240  
catgagacta agatccctca acaatgtata gatctcact acgcaccaac tctaaattga 3300  
cggccgcgtt ggctgttctc ccgatatctg accgaccggt ttgaattttg taggaagcga 3360  
atccctaatt atttaattca atcaacgcac acagcatagt gtacgcgatc agaggacaag 3420  
tcaggctgcc gatcgatgaac gaggaaggga attcgggtgt cgatggagt ctgcaggaag 3480  
ccagagtcca gacggtgcc cagaacttgc cgggtggtaaa gagatcccag agcgagagg 3540

```

ttgagtgggt ggcgttcaag accaacgaca acgcgatggt gaactcgcta gccgggagga 3600
catcggcagtg aagggcgatc cccgcggatg tactggctaa cgcctggagg gtgtcgccgg 3660
aggaggcgag gaggggtgaag ttcaacaggc aggagactca cttggctagc accaggggcc 3720
agtccagggtc gcccgggagg ttgaatgtcg tcaaggaggt gatcaacttg cttatgtaaa 3780
atgtgacgggt gaaataataa cggtaaaata tatgtaataa taataataat aaagccacaa 3840
agtgagaatg aggggaaggg gaaatgtgta atgagccagt agccgggtggt gctaattttg 3900
tatcgtattg tcaataaaatc atgaattttg tggttttttat gtgttttttt aaatcatgaa 3960
ttttaaattt tataaaataa tctccaatcg gaagaacaac attccatata catggatggt 4020
tctttaccca aatctagttc ttgagaggat gaagcatcac cgaacagttc tgcaactatc 4080
cctcaaaagc tttaaaatga acaacaagga acagagcaac gttccaaaga tcccaaacga 4140
aacatattat ctataactat actatattat taattactac tgcccgggat cacaatccct 4200
gaatgattcc tattaactac aagccttggt ggcggcggag aagtgatcgg cgcggcgaga 4260
agcagcggag tcgggagaca ggccttggat gagcagagtc tttacctgcc agggcgtgaa 4320
ggggaagagc ggccttctgg agtaggagtt cagcaagcgg cggttccttg gcggagtaag 4380
cggacgtaag ggtggntgtc gacgtcntcg tttcnggagg cgnattcatg aagggttaaa 4440
gtcanatctg tagctctcga gtgctcaggg agccnaaaga cgttgggaaa ccgtcgnctg 4500
ttggggcatc agtcnccggg gcacgcttcc ctccctgtgc tccanaancn angtanattt 4560
aaaaganatg ggaaattaan taatggnaat nannaggagg attgnaacgg tcnganccgn 4620
angaanagtt tttannggtt taaatactgg gggagtngna gccngccnct ggttccngtg 4680
tagangaaac caagnnccgg gaggnntnca nnnngnnagg agaaaaagga nncatttnan 4740
nangcngagg gacatgaanc ggtacngagc tngngttcan nnancggcgn nnggnagtcc 4800
cnngggaccn ggntggggtn anaagggaan ggaacattng gtngnangga naanaccntt 4860
ttacnattgc ctttgcaggn nngtntnggc ncntnccggg nacatnccgc tgcattgggt 4920
ttggggngcc nanaggnagc cncangggna nncngccncc ttgtncangn cgctnaagtt 4980
cnattgtana tggncgttg

```

<210> 9  
 <211> 96  
 <212> PRT  
 <213> Linum usitatissimum

<400> 9  
 Met Ala Arg Ser Ser Ser Pro Leu Leu Leu Ser Leu Cys Ile Phe Ala  
 1 5 10 15  
 Ile Leu Phe His Ser Ser Leu Gly Arg Gln Gln Phe Gln Gln Gly Asn  
 20 25 30  
 Glu Cys Gln Ile Asp Arg Ile Asp Ala Ser Glu Pro Asp Lys Thr Ile  
 35 40 45  
 Gln Ala Glu Ala Gly Glu Val Trp Asp Gln Asn Arg Gln Gln Phe Gln  
 50 55 60  
 Cys Ala Gly Val Ala Val Val Arg Arg Thr Ile Glu Pro Lys Gly Leu  
 65 70 75 80  
 Leu Leu Pro Phe Tyr Ser Asn Thr Pro Gln Leu Ile Tyr Ile Val Gln  
 85 90 95

<210> 10  
 <211> 85  
 <212> PRT  
 <213> Linum usitatissimum

<400> 10  
 Gly Arg Gly Val Thr Gly Ile Met Phe Pro Xaa Cys Pro Glu Thr Phe  
 1 5 10 15  
 Glu Glu Ser Gln Gln Gln Gly Gln Gln Gly Gln Gly Ser Ser Gln  
 20 25 30  
 Asp Gln His Gln Lys Ile Arg Arg Phe Arg Glu Gly Asp Val Ile Ala  
 35 40 45

Pro Gln Asn Phe Ala Val Val Lys Arg Ser Gln Ser Glu Arg Phe Glu

50		55		60
Trp Val Ala Phe Lys Thr Asn Asp Asn Ala Met Val Asn Ser Leu Ala				
65		70	75	80
Gly Arg Thr Ser Ala Val Arg Ala Ile Pro Ala Asp Val Leu Ala Asn				
	85	90		95
Ala Trp Arg Val Ser Pro Glu Glu Ala Arg Arg Val Lys Phe Asn Arg				
	100	105		110
Gln Glu Thr His Leu Ala Ser Thr Arg Gly Gln Ser Arg Ser Pro Gly				
	115	120	125	
Arg Leu Asn Val Val Lys Glu Val Ile Asn Leu Leu Met				
	130	135	140	

<210> 13  
 <211> 11  
 <212> PRT  
 <213> Linum usitatissimum

<400> 13  
 Gln Gln Gln Gly Gln Gln Gly Gln Gln Gln  
 1 5 10

<210> 14  
 <211> 18  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Primer

<400> 14  
 tccactatgt aggtcata 18

<210> 15  
 <211> 18  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Primer

<400> 15  
 ctttaaggtg tgagagtc 18

<210> 16  
 <211> 15  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Primer

<400> 16  
 aggggtgatc gatta 15

<210> 17  
 <211> 18  
 <212> DNA  
 <213> Artificial Sequence

005280 " 6554960

32

<223> Description of Artificial Sequence: HindIII Site

aagctt

6

<211> 29

<212> DNA

<213> Artificial Sequence

<223> Description of Artificial Sequence: Primer

taggtacctg gcaggtaaag actctgctc

29

<211> 6

<212> DNA


<213> Artificial Sequence

<223> Description of Artificial Sequence: KpnI Site

ggtacc

6

United States Patent & Trademark Office  
Office of Initial Patent Examination -- Scanning Division



Application deficiencies were found during scanning:

☐ Page(s) \_\_\_\_\_ of \_\_\_\_\_ were not present  
for scanning. (Document title)

☐ Page(s) \_\_\_\_\_ of \_\_\_\_\_ were not present  
for scanning. (Document title)

☒ Scanned copy is best available. *Drawings.*